

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau(43) International Publication Date
5 July 2001 (05.07.2001)

PCT

(10) International Publication Number
WO 01/48183 A2

- (51) International Patent Classification?: C12N 15/00
- (21) International Application Number: PCT/EP00/13149
- (22) International Filing Date:
22 December 2000 (22.12.2000)
- (25) Filing Language: English
- (26) Publication Language: English
- (30) Priority Data:
9930691.2 24 December 1999 (24.12.1999) GB
- (71) Applicant (for all designated States except US): DEV-GEN NV [BE/BE]; Technologiepark 9, B-9052 Zwijnaarde (BE).
- (72) Inventors; and
- (75) Inventors/Applicants (for US only): PLAETINCK, Geert [BE/BE]; Pontstraat 16, B-9820 Merelbeke (BE). MORTIER, Katherine [BE/BE]; Paddenhoek 20, B-9830 St.-Martens Latem (BE). LISSENS, Ann [BE/BE]; Tiensesteenweg 137, B-3010 Kessel-Lo (BE). BOGAERT, Thierry [BE/BE]; Wolvendreef 26g, B-8500 Kortrijk (BE).
- (74) Agent: BAYLISS, Geoffrey, Cyril; Boulton Wade Tennant, Verulam Gardens, 70 Gray's Inn Road, London WC1X 8BT (GB).
- (81) Designated States (national): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.
- (84) Designated States (regional): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

Published:

— Without international search report and to be republished upon receipt of that report.

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

WO 01/48183 A2

(54) Title: IMPROVEMENTS RELATING TO DOUBLE-STRANDED RNA INHIBITION

(57) Abstract: There are described ways of improving the efficiency of double stranded RNA inhibition as a method of inhibiting gene expression in nematode worms such as *C. elegans*. In particular, the invention relates to the finding that changes in the genetic background of *C. elegans* result in increased sensitivity to double-stranded RNA inhibition.

IMPROVEMENTS RELATING TO DOUBLE-STRANDED RNA
INHIBITION

5 The present invention is concerned with ways of
improving the efficiency of double stranded RNA
inhibition as a method of inhibiting gene expression
in nematode worms such as *C. elegans*. In particular,
the invention relates to the finding that the
susceptibility of nematode worms such as *C. elegans* to
10 double stranded RNA inhibition is affected by changes
in the genetic background of the worms.

It has recently been described in Nature Vol 391,
pp.806-811, February 98, that introducing double
stranded RNA into a cell results in potent and
15 specific interference with expression of endogenous
genes in the cell, which interference is substantially
more effective than providing either RNA strand
individually as proposed in antisense technology. This
specific reduction of the activity
20 of the gene was also found to occur in the nematode
worm *Caenorhabditis elegans* (*C. elegans*) when the RNA
was introduced into the genome or body cavity of the
worm.

The present inventors have utilized the double
25 stranded RNA inhibition technique and applied it
further to devise novel and inventive methods of (i)
assigning functions to genes or DNA fragments which
have been sequenced in various projects, such as, for
example, the human genome project and which have yet
30 to be accorded a particular function, and (ii)
identifying DNA responsible for conferring a
particular phenotype. Such methods are described in
the applicant's co-pending application number WO
00/01846. Processes for introducing RNA into a living
35 cell, either *in vivo* or *ex vivo*, in order to inhibit
expression of a target gene in that cell are

additionally described in WO 99/32619.

Several different experimental approaches can be used to introduce double-stranded RNA into nematode worms in order to achieve RNA interference *in vivo*.
5 One of the most straightforward approaches is simple injection of double-stranded RNA into a body cavity. A more elegant solution is to feed the nematodes on food organisms, generally bacteria, which express a double stranded RNA of the appropriate sequence,
10 corresponding to a region of the target gene.

The present inventors have now determined that the phenomenon of RNA interference in nematodes following ingestion of food organisms capable of expressing double-stranded RNA is dependent both on
15 the nature of the food organism and on the genetic background of the nematodes themselves. These findings may be exploited to provided improved methods of double-stranded RNA inhibition.

Therefore, according to a first aspect of the present invention there is provided a method of
20 inhibiting expression of a target gene in a nematode worm comprising feeding to said nematode worm a food organism which is capable of producing a double-stranded RNA structure having a nucleotide sequence
25 substantially identical to a portion of said target gene following ingestion of the food organism by the nematode, wherein the nematode has a non wild-type genetic background selected to provide increased sensitivity to RNA interference as compared to wild
30 type.

Caenorhabditis elegans is the preferred nematode worm for use in the method of the invention although the method could be carried out with other nematodes and in particular with other microscopic nematodes,
35 preferably microscopic nematodes belonging to the genus *Caenorhabditis*. As used herein the term "microscopic" nematode encompasses nematodes of

approximately the same size as *C. elegans*, being of the order 1mm long in the adult stage. Microscopic nematodes of this approximate size can easily be grown in the wells of a multi-well plate of the type generally used in the art to perform mid- to high-throughput screening.

It is an essential feature of this aspect of the invention that the nematode has a non wild-type genetic background which confers greater sensitivity to RNA interference phenomena (abbreviated herein to RNAi) as compared to the equivalent wild type nematodes. As illustrated in the accompanying examples, introduction of double-stranded RNA (abbreviated herein to dsRNA) into a non wild-type strain according to the invention results in greater inhibition of expression of the target gene. Depending on the nature of the target gene, this greater level of inhibition may be detectable at the phenotypic level as a more pronounced phenotype.

The nematode having non wild-type genetic background may, advantageously, be a mutant strain. Mutations which have the effect of increasing susceptibility of the nematode to RNAi may, for example, affect the stability of dsRNA or the kinetics of dsRNA turnover within cells of the worm or the rate of uptake of dsRNA synthesised by a food organism. Suitable mutant strains include mutant strains exhibiting knock-out or loss-of-function mutations in one or more genes encoding proteins involved in RNA synthesis, RNA degradation or the regulation of these processes.

In one preferred embodiment, the nematode is a mutant strain, more preferably a mutant *C. elegans*, which exhibits reduced activity of one or more nucleases compared to wild-type. Suitable strains include mutant strains exhibiting knock-out or loss-of-function mutations in one or more genes encoding

nucleases, such as RNases. A particularly preferred example is the *nuc-1* strain. This mutant *C. elegans* strain is known per se in the art.

5 In a second preferred embodiment, the nematode is a mutant strain, more preferably a mutant *C. elegans*, which exhibits increased gut uptake compared to wild-type. Particularly preferred examples of such strains are the so-called *C. elegans* gun mutants described herein. In a still further embodiment, the nematode
10 may be a transgenic worm comprising one or more transgenes which increase gut uptake relative to wild-type.

The term "increased gut uptake" as used herein is taken to mean increased uptake of foreign particles
15 from the gut lumen and may encompass both increased gut permeability and increased gut molecular transport compared to wild-type *C. elegans*.

C. elegans feeds by taking in liquid containing its food (e.g. bacteria). It then spits out the
20 liquid, crushes the food particles and internalises them into the gut lumen. This process is performed by the muscles of the pharynx. The process of taking up liquid and subsequently spitting it out is called pharyngeal pumping. Once the food particles have been
25 internalised via pharyngeal pumping their contents must cross the gut itself in order to reach target sites in the worm. There are multiple factors which effect the uptake of compounds from the gut lumen to the surrounding tissues. These include the action of
30 multi-drug resistance proteins, multi-drug resistance related proteins and the P450 cytochromes as well as other enzymes and mechanisms available for transport of molecules through the gut wall.

C. elegans mutants which exhibit increased uptake
35 of foreign molecules through the gut may be obtained from the *C. elegans* mutant collection at the *C.*

C. elegans Genetic Center, University of Minnesota, St Paul, Minnesota, or may be generated by standard methods. Such methods are described by Anderson in Methods in Cell Biology, Vol 48, "C. *elegans*: Modern biological analysis of an organism" Pages 31 to 58. Several selection rounds of the PCR technique can be performed to select a mutant worm with a deletion in a desired gene. Alternatively, a population of worms could be subjected to random mutagenesis and worms exhibiting the desired characteristic of increased gut uptake selected using a phenotypic screen, such as the dye uptake method described herein.

As an alternative to mutation, transgenic worms may be generated with the appropriate characteristics. Methods of preparing transgenic worms are well known in the art and are particularly described by Craig Mello and Andrew Fire, Methods in Cell Biology, Vol 48, Ed. H.F. Epstein and D.C. Shakes, Academic Press, pages 452-480.

Worms exhibiting the desired characteristics of increased gut uptake can be identified using a test devised by the inventors based on uptake of a marker precursor molecule which is cleaved by the action of enzymes present in the gut lumen to generate a marker molecule which produces a detectable signal, such as fluorescence. A suitable marker precursor molecule is the fluorescent dye precursor BCECF-AM available from Molecular Probes (Europe BV), Netherlands. This dye only becomes fluorescent when cleaved by esterases and maintained at a pH above 6. The pH of the gut lumen is usually 5 or below. Thus, any BCECF-AM taken up through the pharynx into the gut lumen is not fluorescent until cleaved and the cleaved portion has entered the cells surrounding the lumen which are at a higher pH. Thus, this dye is able to quickly identify mutant or otherwise modified worms which have increased gut transport or permeability. There is a

gradual increase in fluorescence in the tissues surrounding the gut while the gut lumen remains dark. The fluorescence can be detected at an excitation wavelength of 485 nm and an emission wavelength of 530 nm.

5
10
15
20
25
30
35

Specific examples of gun mutant strains isolated using this procedure which may be used in the method of the invention are strains bg77, bg84, bg85 and bg86, although it is to be understood that the invention is in no way limited to the use of these specific strains. The *C. elegans* gun mutant strain bg85 was deposited on 23 December 1999 at the BCCM/LMG culture collection, Laboratorium Voor Microbiologie, Universiteit Gent, K. L. Ledeganckstraat 35, B-9000, Gent, Belgium under accession number LMBP 5334CB. The phrase "the bg85 mutation" as used herein refers to the specific mutation(s) present in the bg85 strain which is/are responsible for conferring the gun phenotype.

20
25
30
35

It is also within the scope of the invention to use a non wild-type nematode strain, preferable a *C. elegans* strain, having multiple mutations which affect sensitivity to RNAi. A preferred type of multiple mutant is one having at least one mutation which results in reduced nuclease activity compared to wild type and at least one mutation which results in increased gut uptake compared to wild type. An example of such a mutant is a *C. elegans* strain having the *nuc-1* mutation and at least one further gun mutation. As exemplified herein, double mutants having the *nuc-1* mutation and a gun mutation exhibit enhanced sensitivity to RNAi as compared to either *nuc-1* or gun single mutants.

35

For the avoidance of doubt, where particular characteristics or properties of nematode worms are described herein by relative terms such as "enhanced"

or "increased" or "decreased" this should be taken to mean enhanced, increased or decreased relative to wild-type nematodes. In the case of *C. elegans*, wild-type is defined as the N2 Bristol strain which is well known to workers in the *C. elegans* field and has been extremely well characterised (see Methods in Cell Biology, Volume 48, *Caenorhabditis elegans: Modern biological analysis of an organism*, ed. by Henry F. Epstein and Diane C. Shakes, 1995 Academic Press; *The nematode Caenorhabditis elegans*, ed. by William Wood and the community of *C. elegans* researchers., 1988, Cold Spring Harbor Laboratory Press; *C. elegans II*, ed. by Donald L. Riddle, Thomas Blumenthal, Barbara J. Meyer and James R. Priess, 1997, Cold Spring Harbor Laboratory Press). The N2 strain can be obtained from the *C. elegans* Genetic Center, University of Minnesota, St Paul, Minnesota, USA.

The food organism for use in the above aspect of the invention is preferably a bacterium such as, for example, a strain of *E.coli*. It will, however, be appreciated that any other type of food organism on which nematodes feed and which is capable of producing dsRNA could be used. The food organism may be genetically modified to express a double-stranded RNA of the appropriate sequence, as will be understood with reference to the examples included herein. One convenient way in which this may be achieved in a bacterial food organism is by transforming the bacterium with a vector comprising a promoter or promoters positioned to drive transcription of a DNA sequence to RNA capable of forming a double-stranded structure. Examples of such vectors will be further described below.

The actual step of feeding the food organism to the nematode may be carried out according to procedures known in the art, see WO 00/01846.

Typically the feeding of the food organisms to the nematodes is performed on standard agar plates commonly used for culturing *C. elegans* in the laboratory. However, the step of feeding the food
5 organism to the nematodes may also be carried out in liquid culture, for example in the wells of 96-well microtitre assay plates.

The inventors have further observed that variations in the food organism can result in enhanced
10 *in vivo* RNAi when the food organism is ingested by a nematode worm.

Accordingly, in a further aspect the invention provides a method of inhibiting expression of a target gene in a nematode worm comprising feeding to said
15 nematode worm a food organism capable of producing a double-stranded RNA structure having a nucleotide sequence substantially identical to a portion of said target gene following ingestion of the food organism by the nematode, wherein the food organism carries a
20 modification selected to provide increased expression or persistence of the double-stranded RNA compared to a food organism which does not carry the modification.

The modification present in the food organism can be any modification which results in increased
25 expression of the dsRNA or in increased persistence of the dsRNA. Suitable modifications might include mutations within the bacterial chromosome which affect RNA stability and/or degradation or mutations which have a direct effect on the rate of transcription. In
30 a preferred embodiment, the food organism is an RNase III minus *E. coli* strain, or any other RNase negative strain.

According to a still further aspect of the invention there is provided a method of inhibiting
35 expression of a target gene in a nematode worm comprising introduction of a DNA capable of producing a double-stranded RNA structure having a nucleotide

sequence substantially identical to a portion of said target gene in said nematode, wherein the nematode is one which exhibits increased gut uptake compared to wild type.

5 In addition to exhibiting increased sensitivity to RNAi following feeding with food organisms capable of expressing a dsRNA, nematodes which exhibit increase gut uptake as described herein also show increased uptake of DNA molecules capable of producing
10 double-stranded RNA structures following ingestion into a nematode.

In a preferred embodiment, the DNA is in the form of a vector comprising a promoter or promoters orientated to relative to a sequence of DNA such that
15 they are capable of driving transcription of the said DNA to make RNA capable of forming a double-stranded structure upon binding of an appropriate RNA polymerase to the promoter or promoters.

Several different arrangements of promoters may
20 be used in such a vector. In a first arrangement a DNA fragment corresponding to a region of the target gene is flanked by two opposable polymerase-specific promoters which are preferably identical. Transcription from the opposable promoters produces
25 two complementary RNA strands which can anneal to form an RNA duplex. The plasmid pGN1 described herein is an example of a vector comprising two opposable T7 promoters flanking a multiple cloning site for insertion of a DNA fragment of the appropriate
30 sequence, corresponding to a region of a target gene. pGN8 is an example of a vector derived from pGN1 containing a fragment of the *C. elegans unc-22* gene. In an alternative arrangement, DNA fragments corresponding to a region of the target gene may be
35 placed both in the sense and the antisense orientation downstream of a single promoter. In this case, the sense/antisense fragments are co-transcribed to

generate a single RNA strand which is self-complementary and can therefore form an RNA duplex.

In both of the above arrangements, the polymerase-specific T3, T7 and SP6 promoters, all of which are well known in the art, are useful for driving transcription of the RNA. Expression from these promoters is dependent on expression of the cognate polymerase. Advantageously, the nematode itself may be adapted to express the appropriate polymerase. Expression of the polymerase may be general and constitutive, but could also be regulated under a tissue-specific promoter, an inducible promoter, a temporally regulated promoter or a promoter having a combination of such characteristics. Transgenic *C. elegans* strains harboring a transgene encoding the desired polymerase under the control of an appropriately-regulated promoter can be constructed according to methods known *per se* in the art and described, for example, by Craig Mello and Andrew Fire in *Methods in Cell Biology*, Vol 48, Ed. H. F. Epstein and D. C. Shakes, Academic Press, pp 452-480.

The advantage of adapting the nematode to express the required polymerase is that it is possible to control inhibition of expression of the target gene in a tissue-specific and/or temporally specific manner by placing expression of the polymerase under the control of an appropriately regulated promoter.

Introduction of DNA into nematodes in accordance with the method of the invention can be achieved using a variety of techniques, for example by direct injection into a body cavity or by soaking the worms in a solution containing the DNA. If the DNA is in the form of a vector as described herein, e.g. a plasmid harboring a cloned DNA fragment between two flanking T7 promoters, then dsRNA corresponding to this DNA fragment will be formed in the nematode resulting in down regulation of the corresponding gene. The

introduced DNA can form an extrachromosomal array,
which array might result in a more catalytic knock-out
or reduction of function phenotype. The DNA might also
become integrated into the genome of the nematode,
5 resulting in the same catalytic knock out or reduction
of function phenotype, but which is stably
transmittable.

In each aspect of the invention, the double-
stranded RNA structure may be formed by two separate
10 complementary RNA strands or a single self-
complementary strand, as described above. Inhibition
of target gene expression is sequence-specific in that
only nucleotide sequences corresponding to the duplex
region of the dsRNA structure are targeted for
15 inhibition.

It is preferred to use dsRNA comprising a
nucleotide sequence identical to a portion of the
target gene, although RNA sequences with minor
variations such as insertions, deletions and single
20 base substitutions may also be used and are effective
for inhibition. It will be readily apparent that 100%
sequence identity between the dsRNA and a portion of
the target gene is not absolutely required for
inhibition and the phrase "substantially identical" as
25 used herein is to be interpreted accordingly.
Generally sequences which are substantially identical
will share at least 90%, preferably at least 95% and
more preferably at least 98% nucleic acid sequence
identity. Sequence identity may be conveniently
30 calculated based on an optimal alignment, for example
using the BLAST program accessible at
WWW.ncbi.nlm.nih.gov.

The invention will be further understood with
reference to the following non-limiting Examples,
35 together with the accompanying Figures in which:

Figure 1 is a plasmid map of the vector pGN1

containing opposable T7 promoters flanking a multiple cloning site and an ampicillin resistance marker.

5 Figure 2 is a plasmid map of the vector pGN8 (a genomic fragment of the *C. elegans unc-22* gene cloned in pGN1).

10 Figure 3 is a plasmid map of the vector pGN29 containing two T7 promoters and two T7 terminators flanking *Bst*XI sites. This vector permits cloning of DNA fragments linked to *Bst*XI adaptors.

15 Figure 4 is a plasmid map of the vector pGN39 containing two T7 promoters and two T7 terminators flanking attR recombination sites (based on the Gateway™ cloning system of Life Technologies, Inc).

20 Figure 5 is a plasmid map of the vector pGX22 (a fragment of the *C. elegans* gene C04H5.6 cloned in pGN29).

25 Figure 6 is a plasmid map of the vector pGX52 (a fragment of the *C. elegans* gene K11D9.2b cloned in pGN29).

Figure 7 is a plasmid map of the vector pGX104 (a fragment of the *C. elegans* gene Y57G11C.15 cloned in pGN29).

30 Figure 8 is a plasmid map of the vector pGZ8 (a fragment of the *C. elegans* gene T25G3.2 cloned in pGN39).

35 Figure 9 shows the results of an RNAi experiment in which wild-type (N2) or *nuc-1* strain *C. elegans* in liquid culture were fed with *E. coli* containing the

plasmid pGX22.

5 Figure 10 shows the results of an RNAi experiment in which wild-type (N2) or *nuc-1* strain *C. elegans* in liquid culture were fed with *E. coli* containing the plasmid pGX52.

10 Figure 11 shows the results of an RNAi experiment in which wild-type (N2) or *nuc-1* strain *C. elegans* in liquid culture were fed with *E. coli* containing the plasmid pGXGZ8.

15 Figure 12 shows the results of an RNAi experiment in which wild-type (N2) or *nuc-1* strain *C. elegans* in liquid culture were fed with *E. coli* containing the plasmid pGX104

20

Example 1

Influence of genetic background on the efficiency of RNAi in *C. elegans*.

5 Introduction

Various different *C. elegans* strains were fed with different bacteria, to test the possibility of RNAi by feeding *C. elegans* with bacteria that produce dsRNA. The possibility of DNA delivery and dsRNA delivery has previously been envisaged by using different bacterial strains. In this experiment the importance of the *C. elegans* strain as receptor of the dsRNA is also shown.

For this experiment the following *E. coli* strains were used:

1. MC1061: F-*araD139* Δ (*ara-leu*)7696 *galE15 galK16* Δ (*lac*)X74 *rps1* (*Str^r*) *hsdR2* (*r_k⁻ m_k⁺*) *mcrA mcrB1*
- regular host for various plasmids,
- Wertman et al., (1986) Gene 49:253-262,
- Raleigh et al., (1989) in Current Protocols in Molecular Biology eds. Ausubel et al, Publishing associates and Wiley Interscience; New York. Unit 1.4.
2. B21(DE3): F- *ompT(lon)* *hsdS_B* (*r_B⁻, m_B⁻*; an *E. coli* B strain) with DE3, a λ prophage carrying the T7 RNA polymerase gene.
- regular host for IPTG inducible T7 polymerase expression,
- Studier et al. (1990) Meth. Enzymol. 185:60-89
3. HT115 (DE3): F- *mcrA mcrB* IN(*rrnD-rrnE*) 1 λ -*rnc14::tr10* (DE3 lysogen: *lacUV5* promoter-T7polymerase)
- host for IPTG inducible T7 polymerase

expression,

- RNaseIII-,

- Fire A, Carnegie Institution, Baltimore, MD,
Pers. Comm.

5

For this experiment the following *C. elegans* strains were used:

10

1. *C. elegans* N2: regular WT laboratory strain

15

2. *C. elegans nuc-1(e1393)*: *C. elegans* strain with a reduced endonuclease activity (>95%); condensed chromatin persists after programmed cell death; ingested (bacterial) DNA in the intestinal lumen is not degraded. Several alleles are described: e1392 (strong allele: has been used for the experiments described below); n887 (resembles e1392) and n334 (weaker allele)

20

- Stanfield et al. (1998) East Coast Worm meeting abstract 171,

- Anonymous, Worm Breeder's Gazette 1(1):17b

Hevelone et al. (1988) Biochem. Genet. 26:447-461

- Ellis et al., Worm breeder's Gazette 7(2):44

- Babu, Worm Breeder's gazette 1(2):10

25

- Driscoll, (1996) Brain Pathol. 6:411-425

- Ellis et al., (1991) Genetics 129:79-94

For this experiment the following plasmids were used:

30

pGN1: A vector encoding for ampicillin resistance, harbouring a multiple cloning site between two convergent T7 promoters.

35

pGN8: pGN1 containing a genomic fragment of *unc-22*. Decreased *unc-22* expression via RNAi results in a "twitching" phenotype in *C. elegans*.

Experimental conditions

12-well micro-titer plates were filled with approximately 2 ml of NGM agar per well (1 litre of NGM agar: 15g Agar, 1g peptone, 3g NaCl, 1ml cholesterol solution (5 mg/ml in EtOH), with sterile addition after autoclaving of 9.5 ml 0.1M CaCl₂, 9.5 ml 0.1 ml MgSO₄, 25 ml 1M KH₂PO₄/K₂HPO₄ buffer pH 6 and 5 ml nystatin solution (dissolved 10 mg/ml in 1:1 EtOH:CH₃COONH₄ 7.5 M).

The dried plates were spotted with approximately 50 µl of an overnight culture of bacteria. When IPTG induction was required, 50 µl of a 10 mM stock solution of IPTG was dropped on top of the bacteria lawn, and incubated at 37°C for approximately 4 hours. Individual nematodes at the L4 growth stage were then placed in single wells. In each well 4 nematodes, and the plates were further incubated at 20°C for 6 days to allow offspring to be formed. The F1 offspring of the seeded nematodes were tested for the twitching phenotype.

Results

Table 1: Percentage of the offspring that show the twitching phenotype

	MC1061	N2	<i>nuc-1</i>
5	pGN1	0%	0%
	pGN1 + IPTG	0%	0%
	pGN8	0%	0%
	pGN8 + IPTG	0%	0%
10	BL21 (DE3)		
	pGN1	0%	0%
	pGN1 + IPTG	0%	0%
	pGN8	20% (+)	>90% (++)
	pGN8 + IPTG	20% (+)	>90% (++±)
15	HT115 (DE3)		
	pGN1	0%	0%
	pGN1 + IPTG	0%	0%
	pGN8	50% (+±)	>90% (++)
20	pGN8 + IPTG	80% (++)	>90% (+++)

5: %: percentage twitchers

10: +: weak twitching

15: ++: twitching

20: +++: strong twitching

25

Conclusions

The experiment with *E. coli* MC1061 shows that no twitching could be observed in this experiment.

30 Neither the N2 nematodes nor the *nuc-1* nematodes showed any twitchers. This is to be expected as *E. coli* MC1061 does not produce any T7 RNA polymerase, and hence the *unc-22* fragment cloned in pGN8 is not

expressed as dsRNA.

5 The experiment with *E. coli* strain BL21(DE3) and
nematode strain N2 shows expected results. BL21(DE3)
harbouring plasmid pGN1 does not result in any
twitching as the pGN1 vector is an empty vector. BL21
(DE3) harbouring pGN8 results in the expression of
10 unc-22 dsRNA. When this dsRNA is fed to the N2
nematode (indirectly by feeding with the bacteria that
produce the dsRNA), this results in a twitching
phenotype, indicating that the dsRNA is able to pass
the gut barrier and is able to perform its interfering
activity.

15 Surprisingly the RNAi effect of the unc-22 dsRNA was
even more pronounced in *C. elegans* strain *nuc-1* than
in the wild type N2 strain. Although one may expect
that the *nuc-1* mutation results in the non-degradation
or at least in a slower degradation of DNA, as the
20 NUC-1 protein is known to be involved in DNase
activity, we clearly observe an enhancement of the
RNAi induced phenotype in *C. elegans* with a *nuc-1*
background. The *nuc-1* mutation has not been cloned
yet, but it has been described that the gene is
25 involved in nuclease activity, and more particularly
DNase activity. If the NUC-1 protein is a nuclease, it
may also have activity on nuclease activity on dsRNA,
which would explain the enhanced RNAi phenotype. The
nuc-1 gene product may be a nuclease, or a regulator
30 of nuclease activity. As the mode of action of RNAi is
still not understood, it is also possible that the
NUC-1 protein is interfering in the mode of action of
RNAi. This would explain why a *nuc-1* mutant is more
sensitive to RNAi.

35

The experiment with the *E. coli* strain HT115 (DE3)

confirms the experiments with the BL21(DE3) strain. The RNA interference observed with the *unc-22* dsRNA is even higher. In comparison with strain BL21(DE3) this could be expected, as HT115(DE3) is a RNase III minus strain, and hence is expected to produce larger amounts of dsRNA, resulting in more prominent RNAi. This indicates further that the RNAi observed in this experiment is the result of the dsRNA produced by the bacteria fed to the *C. elegans*. Feeding *C. elegans* *nuc-1* with HT115(DE3) harbouring pGN8 also results in higher RNA interference phenotype than feeding the same bacteria to *C. elegans* wild-type strain N2. Once again this indicates that improved RNAi can be realised using a nuclease negative *C. elegans* and more particularly with a with the *C. elegans nuc-1* (e1392) strain.

Summary

RNA interference can be achieved in *C. elegans* by feeding the worms with bacteria that produce dsRNA. The efficiency of this RNA interference is dependent both on the *E. coli* strain and on the genetic background of the *C. elegans* strain. The higher the level of dsRNA production in the *E. coli*, the more RNAi is observed. This can be realised by using efficient RNA expression systems such as T7 RNA polymerase and RNAase negative strains, such as RNaseIII minus strains. In this example the level of dsRNA production varied: HT115(DE3) > BL21(DE3) > MC1061.

RNA interference is high in *C. elegans* strains that are nuclease negative, or that are influenced in their nuclease activity. This can be realised by using a mutant strain such as *C. elegans nuc-1*. In this example the sensitivity to RNAi varied: *C. elegans nuc-1* >> *C. elegans* N2

Example 2

Improved RNAi by feeding dsRNA producing bacteria in selected *C. elegans* strains-Comparison of the *nuc-1* strain with several mutants which show improved gut uptake. (designated herein 'gun' mutants). Strains
5 bg77, bg78, bg83, bg84, bg85, bg86, bg87, bg88 and bg89 are typical gun mutant *C. elegans* strains isolated using selection for increased gut uptake (gun phenotype) with the marker dye BCECF-AM.
10

Experimental conditions:

- 12-well micro-titer plates were filled with approximately 2ml of NGM agar (containing 1ml/l of ampicillin (100µg/ml) and 5 ml of 100mM stock
15 IPTG) per well
- the dried plates were spotted with 25µl of an overnight culture of bacteria (BL21DE3/HT115DE3) containing the plasmids pGN1 (T7prom-T7prom) or pGN8 (T7prom-unc-22-T7prom)
- 20 - individual nematodes at the L4 growth stage were then placed in single wells, one nematode per well
- the plates were incubated at 20°C for 6 days to allow offspring to be formed
- 25 - the adult F1 offspring of the seeded nematodes were tested for the twitching phenotype

Results:

Table 2:

20°C/6d	pGN1 HT115DE3	pGN8 BL2DE3	pGN8 HT115DE3	
5 N2	0	1	1	
<i>nuc-1</i>	0	1-2	3	
bg77	0	1-2	3	
bg78	0	1	1-2	
bg83	0	1	1	
10 bg84	0	1-2	3	
bg85	0	1	3	
bg86	0	1	3	
bg87	0	1	1	
bg88	0	1	1	
15 bg89	0	1	1	

figure legend:

- 0 = no twitching
- 1 = no to weak phenotype
- 2 = clear phenotype
- 3 = strong phenotype

Conclusions

- bacterial strain HT115(DE3) shows a better RNAi sensitivity than bacterial strain BL21(DE3)
- the *nuc-1 C. elegans* strain is a better strain than the Wild-type N2 strain for RNAi sensitivity
- various gun mutants (improved gut uptake mutants) and more particularly the gun mutant strains bg77, bg84, bg85, bg86 show improved sensitivity to RNAi compared to Wild-type.

A double mutant *C. elegans* strain (*nuc-1/gun*) shows even greater sensitivity to RNAi compared to wild-type:

5 Double mutants were constructed to test the prediction
that *gun/nuc* mutants would even show more enhanced
RNAi sensitivity. As an example, the crossing
strategy with *gun* strain *bg85* is shown, similar
crosses can be conducted with other *gun* strains, such
10 as *bg77*, *bg84* and *bg86*.

P0 cross: *gun(bg85)* x WT males

F1 cross: *nuc-1* x *gun(bg85)/+* males

15

F2 cross: *nuc-1* x *gun(bg85)/+*; *nuc-1/0* males (50%)
nuc-1 x *+/+*; *nuc-1/0* males (50%)

20

F3 single: *gun(bg85)/+*; *nuc-1* hermaphrodites (25%)
+/+; *nuc-1* hermaphrodites (75%)

F4 single: *gun(bg85)*; *nuc-1* (1/4 of every 4th
plate high staining with BCECF)

25

F5 retest: *gun(bg85)*; *nuc-1* (100% progeny of F4
singled high staining with BCECF)

30

35

To select for the *gun* phenotype, the fluorescence precursor BCECF-AM is used (obtainable from Molecular probes). The precursor BCECF-AM is cleaved by esterases present in the gut of the worm to generate the dye BCECF which is fluorescent at pH values above 6. This allows selection for worms that have a *gun* phenotype. BCECF-AM is taken up through the pharynx into the gut lumen and is not fluorescent until it has been cleaved, and the BCECF portion has entered the

cells surrounding the lumen. Wild-type worms will show slower or no increase in BCECF fluorescence.

5 **Example 3**

Improved RNAi feeding in liquid culture using *nuc-1(e1393)* *C. elegans*.

Introduction

10 N2 and *nuc-1* *C. elegans* strains were fed with bacteria producing dsRNAs that give lethal phenotypes via RNAi. For this example RNAi was performed in liquid culture instead of on agar plates. We show here for a number of genes that the RNAi effect is more penetrant using
15 the *nuc-1* strain than the N2 strain, and that RNAi can be performed in liquid.

For this experiment the following *E. coli* strains were used:

- 20
1. HT115 (DE3): F- *mcrA mcrb* IN(*rrnD-rrnE*) 1 λ -*rncl4::tr10* (DE3 lysogen: lacUV5 promoter -T7 polymerase)
- host for IPTG inducible T7 polymerase expression
25 - RNaseIII-
- Fire A, Carnegie Institution, Baltimore, MD, Pers. Comm.

30 For this experiment, following *C. elegans* strains were used:

1. *C. elegans* N2: regular WT laboratory strain
2. *C. elegans nuc-1(e1393)*: *C. elegans* strain with a
35 reduced endonuclease activity (>95%); condensed chromatin persists after programmed cell death;

- 5 ingested (bacterial) DNA in the intestinal lumen is not degraded. Several alleles are described: e1392 (strong allele: has been used for the experiments described below); n887 (resembles e1392) and n334 (weaker allele)
- Stanfield et al. (1998) East Coast Worm meeting abstract 171
 - Anonymous, Worm Breeder's Gazette 1(1):17b
 - Hevelone et al. (1988) Biochem. Genet. 26:447-461
 - 10 - Ellis et al., Worm breeder's Gazette 7(2):44
 - Babu, Worm Breeder's gazette 1(2):10
 - Driscoll, (1996) Brain Pathol. 6:411-425
 - Ellis et al., (1991) Genetics 129:79-94

15

For this experiment, the following plasmids that all give lethal phenotypes in *C. elegans* via RNAi were used:

- 20 pGX22: a vector encoding ampicillin resistance, containing a genomic fragment of cosmid C04H5.6 corresponding to a member of the RNA helicase family.
- 25 pGX52: a vector encoding ampicillin resistance, containing a genomic fragment of cosmid K11D9.2b corresponding to sarco/endoplasmic Ca²⁺ ATPase also known as SERCA.
- 30 pGZ18: a vector encoding ampicillin resistance, containing a genomic fragment of cosmid T25G3.2 corresponding to a chitin like synthase gene.
- 35 pGX104: a vector encoding ampicillin resistance, containing a genomic fragment of cosmid Y57G11C.15 corresponding to sec-61, a transport protein.

Experimental conditions

- 5 - 1 ml overnight cultures of HT115 (DE3) bacteria containing the plasmids pGX22, pGX52, pGZ18 or pGX104 respectively were pelleted and resuspended in S-complete medium, containing 1ml/l of ampicillin (100 µg/ml) and 1ml/l of 1000mM IPTG.
- 10 - 10 µl of this bacterial solution was transferred to a 96-well microtiter plate already filled with 100 µl S-complete containing 1ml/l of ampicillin (100 µg/ml) and 1ml/l of 1000mM IPTG.
- 15 - 3 nematodes at the L1 growth stage of N2 and nuc-1 strain were then placed in single wells, 3 L1's per well. Per experimental set up, 16 wells were used (n=16).
- 20 - the plates were incubated at 25°C for 5 days to allow offspring to be formed.
- the plates were visually checked and the following phenotypes could be scored per individual well:
 - 25 **no effect:** L1's developed to adults and gave normal offspring.
 - no F1 offspring:** L1's developed to adults and gave no offspring.
 - 30 **acute lethal:** original L1 did not mature and died.

Results

- 35 The results of this experiment are illustrated graphically in Figures 9 to 12. Data are expressed as

a percentage of the total (n=16) on the y-axis for both N2 and *nuc-1* strains.

Conclusions

- 5 The following genes were tested in this liquid RNAi assay:
- C04H5.6: an RNA helicase. RNAi of this gene interferes with the generation of offspring.
 - 10 - SERCA: a sarco/endoplasmic Ca²⁺ ATPase. A strong RNAi phenotype causes an acute lethal phenotype. A less penetrant RNAi effect results in loss of offspring.
 - T25G3.2: a chitin like synthase gene. RNAi of this gene causes dead eggs.
 - 15 - *sec-61*: a transport protein. A strong RNAi phenotype causes an acute lethal phenotype. A less penetrant RNAi effect results in loss of offspring.
 - 20 - RNAi can be performed under liquid conditions.

As in the previous examples this set of experiments shows that the *nuc-1 C. elegans* strain is more sensitive to RNAi than the wild-type N2 strain. This is most clear for less penetrant phenotypes such as SERCA and chitin synthase. For strong RNAi phenotypes like the helicase and *Sec-61* the difference between the N2 wild-type strain and the *nuc-1* stain is less pronounced.

Example 4**Cloning of pGX22, pGX52, pGZ18 and pGX104 for RNAi**

A set of primers for each gene was designed on the basis of sequence data available in the publicly accessible *C. elegans* sequence database (Acedb).
5

The cosmid names relate to:

1. C04H5.6=member of RNA helicase
- 10 2. K11D9.2b=SERCA
3. Y57G11C.15=transport protein sec-61
4. T25G3.2=chitin synthase like

The following primer sequences were designed:

- 15 1. C04H5.6F 5'-TGCTCAGAGAGTTTCTCAACGAACC-3'
C04H5.6R 5'-CAATGTTAGTTGCTAGGACCACCTG-3'
2. K11D9.2bF 5'-CAGCCGATCTCCGTCTTGTG-3'
20 K11D9.2bR 5'-CCGAGGGCAAGACAACGAAG-3'
3. Y57G11C.15F 5'-ACCGTGGTACTCTTATGGAGCTCG-3'
Y57G11C.15R 5'-TGCAGTGGATTGGGTCTTCG-3'
- 25 4. T25G3.2F
5'-GGGGACAAGTTTGTACAAAAAAGCAGGCTATGCCAAGTACATGTCGATTGCG-3'
- T25G3.2R
5'-GGGGACCACTTTGTACAAGAAAGCTGGGTGGAGAAGCATTCCGAGAGTTTG-3'
- 30

PCR was performed on genomic DNA of N2 strain *C. elegans* to give PCR products of the following sizes:

- 35 1326bp for C04H5.6
- 1213bp for K11D9.2b

1024bp for Y57G11C.15

1115bp for T25G3.2

5 The PCR fragments of C04H5.6, K11D9.2b and Y57G11C.15 were linked to *Bst*XI adaptors (Invitrogen) and then cloned into the pGN29 vector cut with *Bst*XI. pGN29 contains two T7 promoters and two T7 terminators flanking a cloning site which is adapted for facilitated cloning of PCR fragments, comprising a
10 stuffer DNA flanked by two *Bst*XI sites (see schematic Figure 3). The resulting plasmids were designated pGX22 (C04H5.6), pGX52 (K11D9.2b) and pGX104 (Y57G11C.15).

15 The PCR fragment of T25G3.2 was cloned into pGN39 via recombination sites based on the GATEWAY™ cloning system (Life Technologies, Inc). pGN39 contains two T7 promoters and two T7 terminators flanking a cloning site which facilitates "High Throughput" cloning based
20 on homologous recombination rather than restriction enzyme digestion and ligation. As shown schematically in Figure 4, the cloning site comprises *att*R1 and *att*R2 recombination sites from bacteriophage lambda flanking a gene which is lethal to *E. coli*, in this
25 case the *ccd*B gene. This cloning site is derived from the Gateway™ cloning system commercially available from Life Technologies, Inc. The Gateway™ cloning system has been extensively described by Hartley et al. in WO 96/40724 (PCT/US96/10082).

30

Example 5

Selecting *C. elegans* mutations for increased gut uptake (gun) using marker dye BCECF-AM and *unc-31* as background.

5

The screen was performed in *unc-31(e928)* mutant background, to ensure high amounts of dye in the gut lumen, since *unc-31* mutations show constitutive pharyngeal pumping. The dye (BCECF-AM: 2',7' bis (2-carboxyethyl)-5-(and-6)-carboxyfluorescein, acetoxymethylester), obtained from Molecular Probes, is cleaved by intracellular esterases. Fluorescence accumulates in the gut cells upon passage through the apical gut membrane.

10
15Mutagenesis

- Day 1: *unc-31* L4 staged worms were mutagenised with EMS (final concentration 50 mM) for 4 hours
- Day 2: P0 was divided over several large agar plates
- Day 6: F1's were collected and dropped on large plates. The number of eggs the F1's laid were checked every hour and de F1's were removed when 10-20 eggs per F1 were counted
- Day 10: F2 adults were collected and screened with BCECF-AM. Mutations with increased staining of the gut cells after 15-30 minutes exposure to the dye were selected and singled on small agar plates.

20
25
30

About 50 initial positives gave progeny which was retested with BCECF-AM (2x) and leucine CMB (1x) 9 of the 50 strains were kept (2 strains : 3 times positive, 7 other strains : twice positive)

35

Table 3: Isolation of mutations for increased staining with BCECF-AM

Total P0	Total F1	Total F2	screened chromosomes	number of strains isolated
(counted)	(estimated)	(calculated)	(estimated)	(counted)
2251	55618	222472	100000	9

Outcrossing, backcrossing and double construction

- 10 1. backcrossing *unc-31; gun* --> *unc-31; gun*
- *unc-31; gun* x WT males
 - singled 2x5 WT hermaphrodites F1s (*unc-31/+; gun/+*)
 - singled 50 WT hermaphrodites F2s (1/4 homozygous)
 - select strains segregating 1/4 *unc*
- 15 - stain *unc* strains with BCECF-AM
- from positive strains pick *unc* homozygous
 - retest 100 % *unc* strains with BCECF-AM
 - kept 1 strain (backcrossed)
- 20 2. *unc-31* background was crossed out-->+; *gun*
- *unc-31; gun* x WT males
 - singled 2x5 WT hermaphrodites F1s (*unc-31/+; gun/+*)
 - singled 50 WT hermaphrodites F2s (1/4 homozygous)
 - select strains which did not segregate *unc* F3s
- 25 anymore
- stain non *unc* strains with BCECF-AM
 - 7 positive strains were retested with BCECF-AM and finally 1 was selected and kept (outcrossed)
- 30 3. +; *gun* (1x outcrossed) were 2 times backcrossed-->+; *gun* (3x backcrossed)
- *gun* x WT males
 - WT hermaphrodites x F1 males (*gun/+*)
 - singled 10 WT hermaphrodites F2s (1/2 heterozygous)
- 35 - singled 50 WT hermaphrodites F3s (1/8 homozygous)

- stain strains with BCECF-AM- retested positives with BCECF-AM and finally 1 was selected and kept

- 4. *gun* (3x backcrossed) were crossed with *nuc-1(X)* mutant--> *gun; nuc-1*
- 5 - *gun* x WT males
- *nuc-1* x *gun/+* males
- *nuc-1* x *gun/+; nuc-1/0* or *+/+; nuc-1/0* males
- 10 - singled 10 WT hermaphrodite progeny (*nuc-1* homozygous, ½ heterozygous *gun*)
- singled 40 WT hermaphrodite progeny (1/8 homozygous *gun*)
- stain strains with BCECF-AM
- retested positives with BCECF-AM and finally 1 was selected and kept
- 15

Table 6: Strains derived from *gun* mutations

allele number	<i>unc-31; gun</i>		<i>unc-31; gun</i>		<i>+</i> ; <i>gun</i>			<i>gun; nuc-1</i>
	isolation number	strain number	isolation number	strain number	isolation number	strain number	3x b.c.	from 3x b.c.
bg77	31.4	UG 510	31.4.46.1	UG 556	31.4.34	UG 563	UG 674	UG 777
bg78	37.5	UG 511	37.5.46.4	UG 557	37.5.15	UG 564	UG 675	-
bg83	10.2	UG 543	10.2.11	UG 600	10.2.21	UG 586	UG 676	-
bg84	7.2	UG 544	7.2.10	UG 601	7.2.15	UG 589	UG 677	UG 774
bg85	11.5	UG 545	11.5.29.2	UG 602	2x b.c.	UG 717		UG 775
bg86	42.1	UG 546	42.1.4.5	UG 603	42.1.18	UG 587	UG 678	UG 776
bg87	7.1	UG 547	7.1.8.3	UG 604	7.1.22	UG 585	UG 679	-
bg88	5.3	UG 548	5.3.9	UG 605	5.3.18	UG 584	UG 680	-
bg89	23.4	UG 549	23.4.13.5	UG 606	23.4.3	UG 588	UG 671	-

SEQUENCE LISTING:

SEQ ID NO: 1 complete sequence of pGN1

5 SEQ ID NO: 2 complete sequence of pGN8

SEQ ID NO: 3 complete sequence of pGN29

SEQ ID NO: 4 complete sequence of pGN39

10 SEQ ID NO: 5 complete sequence of pGX22

SEQ ID NO: 6 complete sequence of pGX52

15 SEQ ID NO: 7 complete sequence of pGX104

SEQ ID NO: 8 complete sequence of pGZ8

SEQ ID NO: 9 primer C04H5.6F

20 SEQ ID NO: 10 primer C04H5.6R

SEQ ID NO: 11 primer K11D9.2bF

25 SEQ ID NO: 12 primer K11D9.2bR

SEQ ID NO: 13 primer Y57G11C.15F

SEQ ID NO: 14 primer Y57G11C.15R

30 SEQ ID NO: 15 primer T25G3.2F

SEQ ID NO: 16 primer T25G3.2R

Claims:

1. A method of inhibiting expression of a target gene in a nematode worm comprising feeding to
5 said nematode worm a food organism which is capable of producing a double-stranded RNA structure having a nucleotide sequence substantially identical to a portion of said target gene following ingestion of the food organism by the nematode, wherein the nematode
10 has a non wild-type genetic background selected to provide increased sensitivity to RNA interference as compared to wild type.
2. A method as claimed in claim 1 wherein the
15 nematode is a microscopic nematode.
3. A method as claimed in claim 2 wherein the nematode is from the genus *Caenorhabditis*.
- 20 4. A method as claimed in claim 3 wherein the nematode is *C. elegans*.
5. A method as claimed in any one of claims 1 to 4 wherein the nematode has a mutant genetic
25 background.
6. A method as claimed in claim 5 wherein the nematode is a mutant strain which exhibits reduced activity of one or more nucleases compared to wild
30 type.
7. A method as claimed in claim 6 wherein the nematode is *C. elegans* strain *nuc-1*.
- 35 8. A method as claimed in claim 5 wherein the nematode is a mutant strain which exhibits increased

gut uptake compared to wild type.

9. A method as claimed in claim 8 wherein the nematode is mutant *C. elegans* strain bg85.

5

10. A method as claimed in claim 5 wherein the nematode is a mutant strain having at least one mutation which results in reduced nuclease activity compared to wild type and at least one mutation which results in increased gut uptake compared to wild type.

10

11. A method as claimed in claim 10 wherein the nematode is a mutant *C. elegans* strain having the *nuc-1* mutation and the bg85 mutation.

15

12. A method as claimed in any one of the preceding claims wherein the food organism has been engineered to express a double-stranded RNA.

20

13. A method as claimed in any one of the preceding claims wherein the food organism is a bacterium.

25

14. A method as claimed in claim 13 wherein the food organism is *E. coli*.

30

15. A method as claimed in any one of the preceding claims wherein the food organism has been genetically modified to express a double-stranded RNA having a nucleotide sequence substantially identical to a portion of said target gene.

35

16. A method as claimed in claim 15 wherein the food organism contains a DNA vector, the vector comprising a promoter or promoters orientated relative to a DNA sequence such that they are capable of

initiating transcription of said DNA sequence to RNA capable of forming a double-stranded structure upon binding of an appropriate RNA polymerase to said promoter or promoters.

5

17. A method as claimed in claim 25 wherein the vector comprises two promoters flanking the DNA sequence.

10

18. A method as claimed in claim 26 wherein the two promoters are identical.

15

19. A method as claimed in claim 25 wherein the vector comprises a single promoter and further comprises said DNA sequence in a sense and an antisense orientation relative to said promoter.

20

20. A method as claimed in any one of claims 16 to 20 wherein the nematode or the food organism is adapted to express an RNA polymerase capable of initiating transcription from said promoter or promoters.

25

21. A method as claimed in any one of claims 16 to 20 wherein the RNA polymerase is T7, T3 or SP6 polymerase.

30

22. A method as claimed in any one of claims 1 to 21 wherein the step of feeding said food organism to said nematode worm is carried out in liquid culture.

35

23. A method of inhibiting expression of a target gene in a nematode worm comprising feeding to said nematode worm a food organism capable of producing a double-stranded RNA structure having a nucleotide sequence substantially identical to a

nematode is *C. elegans*.

5 32. A method as claimed in any one of claims 28 to 31 wherein the nematode has a mutant genetic background.

33. A method as claimed in claim 32 wherein the nematode is mutant *C. elegans* strain bg85.

10 34. A method as claimed in any one of claims 28 to 33 wherein the DNA capable of producing a double-stranded RNA structure is a vector comprising a promoter or promoters orientated relative to a DNA sequence such that they are capable of initiating
15 transcription of said DNA sequence to RNA capable of forming a double-stranded structure upon binding of an appropriate RNA polymerase to said promoter or promoters.

20 35. A method as claimed in claim 34 wherein the vector comprises two promoters flanking the DNA sequence.

25 36. A method as claimed in claim 35 wherein the two promoters are identical.

30 37. A method as claimed in claim 34 wherein the vector comprises a single promoter and further comprises said DNA sequence in a sense and an antisense orientation relative to said promoter.

35 38. A method as claimed in any one of claims 34 to 37 wherein the nematode is adapted to express an RNA polymerase capable of initiating transcription from said promoter or promoters.

39. A method as claimed in any one of claims 34

portion of said target gene following ingestion of the food organism by the nematode, wherein the food organism carries a modification selected to provide increased expression or persistence of the doubled-stranded RNA compared to a food organism which does not carry the modification.

24. A method as claimed in claim 23 wherein the food organism is a bacterium.

25. A method as claimed in claim 24 wherein the bacterium is an *E. coli* strain.

26. A method as claimed in claim 25 wherein the *E. coli* strain is an RNase III minus strain or any other RNase negative strain.

27. A method as claimed in any one of claims 23 to 26 wherein the step of feeding said food organism to said nematode worm is carried out in liquid culture.

28. A method of inhibiting expression of a target gene in a nematode worm comprising introduction of a DNA capable of producing a double-stranded RNA structure having a nucleotide sequence substantially identical to a portion of said target gene in said nematode, wherein the nematode is one which exhibits increased gut uptake compared to wild type.

29. A method as claimed in claim 28 wherein the nematode is a microscopic nematode.

30. A method as claimed in claim 29 wherein the nematode is from the genus *Caenorhabditis*.

31. A method as claimed in claim 30 wherein the

to 38 wherein the RNA polymerase is T7, T3 or SP6 polymerase.

FIG. 1.




FIG. 2.




FIG. 3.




FIG. 4.




FIG. 5.




FIG. 6.




FIG. 7.




FIG. 8.




FIG. 9.




FIG. 10.




FIG. 11.





FIG. 12.



1
SEQUENCE LISTING

<110> DEVGEN NV

<120> IMPROVEMENTS RELATING TO DOUBLE-STRANDED RNA INHIBITION

<130> SCB/53711/001

<140>

<141>

<160> 14

<170> PatentIn Ver. 2.0

<210> 1

<211> 3216

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Plasmid pGN1

<400> 1

```

gagtgcacca tatgcggtgt gaaataccgc acagatgcgt aaggagaaaa taccgcatca 60
ggcgaaattg taaacgtaa tattttgtta aaattcgcgt taaatatttg ttaaatacagc 120
tcatttttta accaatagggc cgaaatcggc aaaatccctt ataatcaaaa agaatagacc 180
gagatagggg tgagtgttgt tccagtttgg aacaagagtc cactattaaa gaacgtggac 240
tccaacgtca aaggcgaaa aaccgtctat cagggcgatg gccactacg tgaacatca 300
cccaaatcaa gtttttggc gtcgaggtgc cgtaaagctc taaatcgaaa ccctaaaggg 360
agccccgat ttagagcttg acgggaaaag cggcgaaacg tggcgagaaa ggaagggaag 420
aaagcgaaag gagcgggccc tagggcgctg gcaagtgtag cggtcacgct gcgcgtaacc 480
accacacccg ccgcgcttaa tgcgccgcta cagggcgctt ccattcgcca ttcaggctgc 540
gcaactgttg ggaagggcga tcggtgcggg cctcttcgct attacgccag ctggcgaaaag 600
ggggatgtgc tgcaaggcga ttaagtggg taacgccagg gttttccag tcacgacgtt 660
gtaaacgacg gccagtgaa ttgtaatac actcactata gggcgaattc gagctcggta 720
cccggggatc ctctagagtc gaaagcttct cgccctatag tgagtcgtat tacagcttga 780
gtattctata gtgtcaccta aatagcttgg cgtaatcatg gtcatagctg tttcctgtgt 840
gaaattgtta tccgctcaca attccacaca acatacagc cggaagcata aagtgtaaaag 900
cctgggggtg ctaatgagtg agctaactca cattaattgc gttgcgctca ctgcccgtt 960
tccagtcggg aaacctgtcg tgccagctgc attaatgaat cggccaacgc gcggggagag 1020
gcggtttgcg tattgggccc tcttccgctt cctcgctcac tgactcgctg cgctcggctg 1080
ttcggctgcg gcgagcggta tcagctcaact caaaggcggg aatacgggta tccacagaat 1140
caggggataa cgcaggaaa aacatgtgag caaaaggcca gcaaaaggcc aggaaccgta 1200
aaaaggccgc gttgctggcg ttttcgata ggctccgccc ccctgacgag catcacaanaa 1260
atcgacgctc aagtcagagg tggcgaaaacc cgacaggact ataaagatac caggcgtttc 1320
cccctggaag ctccctcgtg cgctctcctg ttccgaccct gccgcttacc ggatacctgt 1380
ccgcctttct ccttcggga agcgtggcgc tttctcatag ctacgctgt aggtatetca 1440
gttcggtgta ggtcgttcgc tccaagctgg gctgtgtgca cgaaccccc gttcagcccg 1500
accgctgccc cttatccggt aactatcgtc ttgagtcaa cccggtaaga cacgacttat 1560
cgccactggc agcagccact ggtaacagga ttagcagagc gaggtatgta ggcggtgcta 1620
cagagttctt gaagtgggtg cctaactacg gctacactag aaggacagta tttgggtatct 1680
gcgctctgct gaagccagtt accttcgaaa aaagagttgg tagctcttga tccggcaaac 1740
aaaccaccgc tggtagcggg gttttttttg tttgcaagca gcagattacg gcgagaaaaa 1800
aaggatctca agaagatcct ttgatctttt ctacggggtc tgacgctcag tggaacgaaa 1860
actcacgtta agggattttg gtcgatgagat tatcaaaaag gatcttcacc tagatccttt 1920
taaattaaaa atgaagtttt aatcaatct aaagtatata tgagtaaact tggctctgaca 1980

```

gttaccaatg	cttaatcagt	gaggcaccta	tctcagcgat	ctgtctat	cgttcatcca	2040
tagttgcctg	actccccgtc	gtgtagataa	ctacgatagc	ggagggctta	ccatctggcc	2100
ccagtgctgc	aatgataaccg	cgagaccac	gctcaccggc	tccagattta	tcagcaataa	2160
accagccagc	cggaagggcc	gagcgcagaa	gtggctctgc	aactttatcc	gectccatcc	2220
agtctattaa	ttgttgccgg	gaagctagag	taagtagttc	gccagttaat	agtttgcgca	2280
acgttgttgg	cattgctaca	ggcatcgtag	tgtcagctc	gtcgtttggt	atggcttcat	2340
tcagctccgg	ttcccaacga	tcaaggcgag	ttacatgatc	ccccatggtg	tgcaaaaaag	2400
cggttagctc	cttcggtcct	ccgatcgtag	tcagaagtaa	gttggccgca	gtgttatcac	2460
tcattggttat	ggcagcactg	cataattctc	ttactgtcat	gccatccgta	agatgctttt	2520
ctgtgactgg	tgagtactca	accaagtcac	tctgagaata	ccgcgcccgg	cgaccgagtt	2580
gctcttgccc	ggcgtcaata	cgggataata	gtgtatgaca	tagcagaact	ttaaaagtgc	2640
tcattcattgg	aaaacgttct	tcggggcgaa	aactctcaag	gatcttaccg	ctggtgagat	2700
ccagttcogat	gtaaccact	cgtagcacca	actgatcttc	agcatctttt	actttcacca	2760
gcgtttctgg	gtgagcaaaa	acaggaagc	aaaatgccgc	aaaaaaggga	ataaggcgca	2820
cacggaaatg	ttgaatactc	atactcttcc	tttttcaata	ttattgaagc	atztatcagg	2880
gttattgtct	catgagcggg	tacatatttg	aatgtattta	gaaaaataaa	caaatagggg	2940
ttccgcgcac	atttcccoga	aaagtgccac	ctgacgtcta	agaaaccatt	attatcatga	3000
cattaacctta	taaaaatagg	cgatcacgca	ggccctttcg	tctcgcgctg	ttcgggtgatg	3060
acggtgaaaa	cctctgacac	atgcagctcc	cggagacggt	cacagcttgt	ctgtaagcgg	3120
atgccgggag	cagacaagcc	cgtagggcgg	cgtagcgggg	tgtagggcggg	tgtaggggct	3180
ggcttaacta	tgcgcatca	gagcagattg	tactga			3216

<210> 2

<211> 4620

<212> DNA


<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Plasmid pGN8

<400> 2

gatccgaatc	tccatgtctg	ttaacagcct	tgacacggaa	tttatattca	tgcccttgag	60
tcaaatcgtc	aacgtggaag	ttggtatcct	tgctctctcc	gcaagcagtc	catctgccag	120
tggcagcatc	ttgcttttca	atgacatagt	gactgatttc	agctcctcca	tcattctctg	180
gttccttcca	tgcaagatca	catccatcct	tgacaatatt	agtgacatcg	agaggtccac	240
gtgggcttgc	tgtagatca	agaacagtaa	ccttcacttc	agcagtgta	gttcattct	300
cgttctctgc	cttgatgata	taggttctctg	tatccgaacg	caaagctctc	ttcacatgga	360
athtagtctt	gccgtcttca	ttgttcaact	tcatacagtc	atcagattcg	actggtggtc	420
cttcgaaagt	ccaagtaatt	gttggagttg	gttcaccact	gactggaatg	ttcaatgaga	480
agtcctgtcc	agccttgacc	ttgatttctt	gaatcgagtt	acgatcgatg	actggtggaa	540
ctataattta	attcaatgat	tattagtaat	tgatttagac	tcttaccatt	tctagccttt	600
gcaacagctg	atgctgaatc	agatggatct	cccaatcctg	ccttgttctt	ggcacggatt	660
ctgaattcgt	actttgatcc	ttccttgaga	tttccaacag	tagcattcgt	ttgtccagct	720
ggaacatgag	caacgtcatt	ccagaatggc	gagaactcgt	ccttcacttc	aacaacgta	780
tcctcgattg	gagcaccacc	gtcgtttgct	ggtggcttcc	attcaaggtc	aacatgatcc	840
ttatcccaat	cagtaatttc	aggagcattt	gtctttcctg	gcttgtaaaa	tggatctttg	900
gcaagtgtgg	ttccgaaggt	ctccaatgga	tcggactctc	cctcagcatt	gacggcagcg	960
acacggaact	gaaaatcaaa	atggtgtagg	caattgagtt	caagattaaa	aaattctcac	1020
tttatattca	tgtccaggaa	taagaccgtc	aacaacagct	gtagtcttat	ctccagcgac	1080
ctttgcagct	ggaaccatc	ttccacttgc	agtatcgta	ttttcgatca	catagttttc	1140
aattggaata	cctccatcat	catctggtgc	acgcaattc	aaagtacat	gatcaccatg	1200
aacatcggaa	acatctaag	gaccatttgg	agaagttggc	ttgtctgaaa	atttaaaata	1260
taaccaaat	aattgaagaa	aactaatgct	caccaataac	attgatctta	acagttgctt	1320
catcttctcc	atgttcattg	acagctttga	tagtgaaagt	tccactgtct	ccaggttcca	1380
tttgcttca	aaccagcttt	gattggtatt	ctgggttatc	aagcttctcg	ccctatagtg	1440
agtcgtatta	cagcttgagt	attctatagt	gtcacctaaa	tagcttggcg	taatcatggt	1500
catagctggt	tcctgtgtga	aattggtatc	cgctcaaat	tccacacaac	atagcagccg	1560
gaagcataaa	gtgtaaagcc	tggggtgcct	aatgagtgag	ctaactcaca	ttaattgctg	1620



<400> 3

tgctcagaga	gtttctcaac	gaacccgatt	tggctagtta	taggtaattt	ttagaacatt	60
tacaaaaaca	gcaaaaaaac	caaacattca	ggatttttgt	ttttaattaa	gaaaaaaatc	120
gatcgctcct	aaattttaat	caatacttcg	aataaaccca	aaaaaaaacg	aaaaaaaatc	180
ctgtttccag	tgtaatgatg	attgacgagg	ctcacgaacg	tactctacac	acggatattc	240
tattcggttt	agtcaaaagat	attgcaagat	tccgaaagga	tttgaagctt	ctcatctctt	300
ctgcaacact	tgacgctgaa	aagttctcca	gtttcttcga	cgacgctccg	atthttccgaa	360
ttccgggacg	cagattcccg	gtggacattt	actatacaca	ggctcccgaa	gcggtactacg	420
tcgacgcggc	tatcgtcaca	attatgcaga	ttcacttgac	ccagccactt	cccggcgata	480
ttttggatt	tctgactggt	caggaagaaa	tcgaaactgt	acaggaagca	cttatggaac	540
ggtcgaaagc	actgggatcg	aagattaagg	agcttattcc	gctgcccgtt	tatgccaatt	600
tgcccagtga	tttgacggcg	aagattttcg	agccaacgcc	gaaagatgcy	agaaaggtag	660
atthttctta	caaatttttt	ccaaaaaaa	atccgagaaa	aatctacaaa	atthtcaggca	720
aaaaactggtt	cattttattc	ctaactagtt	tttagcaaa	cgthtagatt	taacaaaatt	780
gaacaaattt	gaagthttcc	aatthaaaa	ataaatgtht	cggaaagtht	attgaaaaat	840
ctgaaattgc	tatctctcgc	tatctgcaaa	aaaaacactt	taaaaaatgc	tctgthcttt	900
gaaaatttct	aaactgaaaa	atthgaaatt	tctgaaaatt	gtgataattt	tataaaattt	960
tatagaaaat	gtaagcattc	cagaaaaata	tcaaaaattt	cgagaaaatt	ctgaaaaaat	1020
ccagaaatat	taacagaaaa	aaaatcttht	gaaacatctg	aaaatthaaa	taaatthgat	1080
ttacatttht	ttthttggga	tttctthaaa	atcactatga	atthaccact	aaatthtttg	1140
caaaaaatta	ttthtttaat	ttcaaaagaaa	aagcaagaaa	ttthaaaata	tcaaaaagtc	1200
caaatttggt	tcggtgaatt	tttaaaataa	cattthcaag	ataatthtaa	gthaatcaaa	1260
acattccaag	cattthctagt	ttcccaaatt	tctctaaatt	tcaggtggtc	ctagcaacta	1320
acattgccag	cacaatggat	ctcgagggat	cttccatacc	taccagthct	gcgcctgcag	1380
gtcgcggccg	cgactctcta	gacgcgtaag	cttactagca	taacccttg	gggcctctaa	1440
acgggtcttg	aggggtthtt	tgagcttctc	gccctatagt	gagtcgtatt	acagcttgag	1500
tattctatag	tgtcacctaa	atagcttggc	gtaatcatgg	tcatagctgt	ttcctgtgtg	1560
aaattgtht	ccgctcacia	ttccacacia	catacgagcc	ggaagcataa	agtgtaaagc	1620
ctgggtgccc	taatgagtga	gctaactcac	attaatgcy	ttgcgctcac	tgcccgtatt	1680
ccagtcggga	aacctgtcgt	gccagctgca	ttaatgaatc	ggccaacgcy	cggggagagg	1740
cggthtgctg	attgggcgct	cttccgcttc	ctcgcctcact	gactcgcctgc	gctcggctcgt	1800
tcggctgcyg	cgagcgggat	cagctcactc	aaaggcggta	atagcgttat	ccacagaatc	1860
aggggataac	gcaggaagaa	acatgtgagc	aaaaggccag	caaaaggcca	ggaaccgtaa	1920
aaaggccgcy	ttgctggcgt	ttthcgatag	gctccgccc	cctgacgagc	atcacaaaa	1980
tcgacgctca	agtcagaggt	ggcgaaacc	gacaggacta	taaagatacc	aggcgtthtc	2040
ccctggaaag	tccctcgtgc	gctctcctgt	tccgaccctg	ccgcttaccg	gatacctgtc	2100
cgctthctc	ccttcgggaa	cgctggcgct	ttctcatagc	tcacgctgta	ggtatctcag	2160
ttcgggtgtag	gtcgttcgct	ccaagctggg	ctgtgtgcac	gaacccccg	ttcagcccga	2220
ccgctgcgcc	ttatccggta	actatcgtct	tgagtccaac	ccggtaagac	acgacttatc	2280
gccactggca	gcagccactg	gtaacaggat	tagcagagcy	aggatgtag	gcggtgctac	2340
agagthcttg	aagtgggtgc	gtaactacgg	ctacactaga	aggacagtat	thggtatctg	2400
cgctctgctg	aagccagtha	ccttcggaaa	aagagthggt	agctcttgat	ccggcaaaaa	2460
aaccaccgct	ggtagcgggt	gthttthtgt	thgcaagcag	cagattacgc	gcagaaaaaa	2520
aggatctcaa	gaagatcctt	tgatctthtc	tacgggtct	gacgctcagt	ggaacgaaaa	2580
ctcacgttaa	gggattthtg	tcatgagatt	atcaaaaagg	atcttcacct	agatccttht	2640
aaatthaaaa	tgaagthtth	aatcaatcta	aagtataat	gagtaaaact	ggtctgacag	2700
ttaccaatgc	thaatcagtg	aggcacctat	ctcagcgatc	tgtctatthc	gthcatccat	2760
agthgcctga	ctccccgctc	tgtagataac	tacgatacgg	gagggcttac	catctggccc	2820
cagtgctgca	atgataccgc	gagaccacgc	ctcaccgctc	ccagatthtat	cagcaataaa	2880
ccagccagcc	ggaagggccg	agcgcagaa	tggtcctgca	actthtatccg	cctccatcca	2940
gtctattht	tgttgccggg	aagctagagt	aagtthtctg	ccagthtaata	gthtgccgaa	3000
cgthgtggc	attgctacag	gcacgtggt	gtcacgctc	tcgthtggtg	thggtctcatt	3060
cagctccggt	tccaacgat	caaggcaggt	tacatgatcc	ccatgthgt	gcaaaaagtc	3120
ggttagctcc	ttcggctctc	cgatcgtgt	cagaagtaag	thggccgcag	tgttatcact	3180
catggttatg	gcagcactgc	ataatthctc	tactgtcatg	ccatccgtaa	gatgctthtc	3240
tgtgactggt	gagthactca	ccaagtcatt	ctgagaatac	cgcccccggc	gaccgagthg	3300
ctcttgccc	gcgtcaatac	gggataatag	tgtatgacat	agcagaactt	taaaagthct	3360

catcattgga	aaacgttctt	oggggcgaaa	actctcaagg	atcttaccgc	tgttgagatc	3420
cagttcogatg	taaccactc	gtgcacccaa	ctgatcttca	gcatcttcta	ctttcaccag	3480
cgtttctggg	tgagcaaaaa	caggaaggca	aaatgccgca	aaaaagggaa	taagggcgac	3540
acggaaatgt	tgaatactca	tactcttct	ttttcaatat	tattgaagca	tttatcaggg	3600
ttattgtctc	atgagcggat	acatatttga	atgtattttag	aaaaataaac	aaataggggt	3660
tccgcgcaca	tttccccgaa	aagtgccacc	tgacgtctaa	gaaaccatta	ttatcatgac	3720
attaacctat	aaaaatagc	gtatcacgag	gccctttcgt	ctcgcgcggt	tccggtgatga	3780
cggtgaaaac	ctctgacaca	tgacgtctcc	ggagacggtc	acagcttgtc	tgtaacggga	3840
tgccgggagc	agacaagccc	gtcagggcgc	gtcagcgggt	gttggcgggt	gtcggggctg	3900
gcttaactat	goggcatcag	agcagattgt	actgagagtg	caccatagtc	ggtgtgaaat	3960
accgcacaga	tgcgtaagga	gaaaataacc	catcaggcga	aattgtaaac	gttaatat	4020
tgtaaaaatt	cgcgtaaat	atgtgttaaa	tcagctcatt	ttttaaccaa	taggccgaaa	4080
tcggcaaaaat	cccttataaa	tcaaaagaat	agaccgagat	agggttgagt	gttggtccag	4140
tttgaacaa	gagtccacta	ttaaagaacg	tggtactcaa	cgtaaaagg	cgaaaaaccg	4200
tctatcaggg	cgatggcca	ctacgtgaac	catcacccaa	atcaagtttt	ttgcggtcga	4260
ggtgccgtaa	agctctaaat	cggaaacccta	aagggagccc	ccgatttaga	gcttgacggg	4320
gaaagccggc	gaacgtggcg	agaaaggaag	ggaagaaagc	gaaaggagcg	ggcgctaggg	4380
cgctggcaag	tgtagcggtc	acgctgcgcg	taaccaccac	acccgcgcgc	cttaatgcgc	4440
cgctacaggg	cgctccatt	cgccattcag	gctgcgcaac	tgttgggaa	ggcgatcgg	4500
gcgggcctct	tcgctattac	gccagctggc	gaaaggggga	tgtgctgcaa	ggcgattaag	4560
ttgggtaacg	ccagggttt	cccagtcacg	acgttgtaaa	acgacggcca	gtgaattgta	4620
atagactca	ctatagggcg	aattcaaaaa	accctcaag	acccgtttag	aggcccaag	4680
gggttatgct	agtgaattct	gcagggatcc	cggggatcct	ctagagatcc	ctcgacctcg	4740
agatccattg	tgctgg					4756

<210> 4
 <211> 4643
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence: Plasmid pGX52

<400> 4						
gagtgaccca	tatgcggtgt	gaaataccgc	acagatgcgt	aaggagaaaa	taccgcatca	60
ggcgaaattg	taaacgttaa	tattttgtta	aaattcgcgt	taaataattg	ttaaatcagc	120
tcatttttta	accaataggc	cgaaatcggc	aaaatccctt	ataaatcaaa	agaatagacc	180
gagataggg	tgagtgtgt	tccagtttgg	aacaagagtc	cactattaaa	gaacgtggac	240
tccaacgtca	aagggcgaaa	aaccgtctat	cagggcgatg	gccactacg	tgaaccatca	300
cccaaatcaa	gttttttgcg	gtcagagtg	cgtaaagctc	taaactcgaa	ccctaaagg	360
agccccgat	ttagagcttg	acggggaaag	ccggcgaacg	tggcgagaaa	ggaagggaac	420
aaagcgaag	gagcgggcgc	tagggcgctg	gcaagtgtag	cggtcacgct	gcgcgtaacc	480
accacacccg	ccgcgcttaa	tgcgccccta	cagggcgctg	ccattcgcca	ttcaggtg	540
gcaactggtg	ggaagggcga	tccggtcggg	cctcttcgct	attacgccag	ctggcgaaag	600
ggggatgtgc	tgcaaggoga	ttaagttggg	taacgccagg	gttttcccag	tcacgacgtt	660
gtaaaacgac	ggccagtga	ttgtaatacg	actcactata	gggcgaattc	aaaaaacccc	720
tcaagacccg	tttagaggcc	ccaaggggtt	atgctagtga	attctgcagg	gtaccgggg	780
atcctctaga	gatccctcga	cctcgagatc	cattgtgctg	gcagccgac	tccgtcttgt	840
gaagatctac	tccaccacca	tccgtatcga	tcagtcctac	ctcaccggag	aatctgtg	900
tgttatcaag	cacaccgact	ctgtgccaga	tccacgcgct	gtaaacagg	acaagaagaa	960
ttgtctgttc	tcgggaacca	atgtcgcac	tggaaggct	cgtggaatcg	tcttcggaac	1020
cggattgacc	actgaaatcg	gaaagatccg	taccgaaatg	gctgagaccg	agaatgagaa	1080
gacaccactt	caacagaagt	tgacgaatt	cggagagcaa	ctttccaagg	ttatctctgt	1140
tatttgcgtt	gctgtttggg	ctatcaacat	tggaacttc	aacgatccag	ctcaggtgg	1200
atcatgggtt	aagggagcaa	tctactact	caaaatcgcc	gttgctcttg	ccgtcgtg	1260
tattccagaa	ggacttccag	ctgtcatcac	cacgtgcctt	gcctcggaa	ctcgcggtat	1320
ggccaagaag	aacgctattg	taagatccct	tccatccgtc	gaaactcttg	gatgcacatc	1380
tgttatctgc	tctgacaaga	ctggaactct	caccaccaac	cagatgtctg	tgtaaacgat	1440

gttcatcgct	ggacaagctt	ctggagacaa	catcaacttc	accgagttcg	ccatctccgg	1500
atccacctac	gagccagtcg	gaaaggtttc	caccaatgga	cgtgaaatca	accagctgc	1560
tggagaattc	gaatcactca	ccgagttggc	catgatctgc	gctatgtgca	atgattcatc	1620
tgttgattac	aatgagacca	agaagatcta	cgagaaagtc	ggagaagcca	ctgaaactgc	1680
tcttatcgtt	cttgctgaga	agatgaatgt	tttcggaacc	tcgaaagccg	gactttcacc	1740
aaaggagctc	ggaggagttt	gcaaccgtgt	catccaacaa	aatggaaga	aggagttcac	1800
actcgagttc	tcccgtgatc	gtaaatccat	gtccgcctac	tgcttcccag	cttccggagg	1860
atctggagcc	aagatgttcg	tgaagggagc	cccagaagga	gttctcgga	gatgcacca	1920
cgtcagagtt	aacggacaaa	aggttccact	cacctctgcc	atgactcaga	agattgttga	1980
ccaatgcgtg	caatacggaa	ccggaagaga	taccttctgt	tgtcttgccc	tcggccagca	2040
caatggatct	cgagggatct	tccataccta	ccagttctgc	gcctgcaggt	cgcgcccgcg	2100
actctctaga	cgcgtaagct	tactagcata	acccttggg	gcctctaaac	gggtcttgag	2160
gggttttttg	agcttctcgc	cctatagtga	gtcgtattac	agcttgagta	ttctatagtg	2220
tcacctaaat	agcttggcgt	aatcatggtc	atagctgttt	cctgtgtgaa	attgttatcc	2280
gctcacaatt	ccacacaaca	tacgagccgg	aagcataaag	tgtaaagcct	ggggtgccta	2340
atgagtgagc	taactcacat	taattgcgtt	gcgctcactg	cccgtttcc	agtcgggaaa	2400
cctgtcgtgc	cagctgcatt	aatgaatcgg	ccaacgcgcg	gggagaggcg	gtttgcgat	2460
tgggcgctct	tccgcttcc	cgctcactga	ctcgcgtcgc	tcggtcgttc	ggctgcgcg	2520
agcggatca	gctcactcaa	aggcggtaat	acggttatcc	acagaatcag	gggataaccg	2580
aggaaagaac	atgtgagcaa	aaggccagca	aaaggccagg	aaccgtaaaa	aggccgcgct	2640
gctggcgctt	ttcgataggc	tccgcccc	tgacgagcat	cacaaaaatc	gacgctcaag	2700
tcagaggtgg	cgaaaccoga	caggactata	aagataccag	gcgtttcccc	ctggaagctc	2760
ctcgtgcgc	tctcctgttc	cgaccctgcc	gcttacggga	tacctgtccg	cctttctccc	2820
ttcgggaagc	tgggcgcttt	ctcatagctc	acgctgtagg	tatctcagtt	cgggtgtagt	2880
cgttcgtctc	aagctgggct	gtgtgcacga	acccccgtt	cagcccgacc	gctgcgcctt	2940
atccggtaac	tatcgtcttg	agtccaacc	ggtaagacac	gacttatcgc	cactggcagc	3000
agccactggt	aacaggatta	gcagagcgag	gtatgtaggc	ggtgctacag	agttcctgaa	3060
gtggtggcct	aactacggct	acactagaag	gacagtattt	ggtatctgcg	ctctgctgaa	3120
gccagttacc	ttcgaaaaaa	gagttggtag	ctcttgatcc	ggcaaaaaaa	ccaccgctgg	3180
tagcggtggt	tttttgttt	gcaagcagca	gattacgcgc	agaaaaaaag	gatctcaaga	3240
agatcctttg	atctttcta	cggggctctga	cgctcagttg	aacgaaaaact	cacgttaagg	3300
gattttggtc	atgagattat	caaaaaggat	cttcaactag	atccttttaa	attaaaaatg	3360
aagttttaaa	tcaatctaaa	gtatatatga	gtaaaacttg	tctgacagtt	accaatgctt	3420
aatcagtgag	gcacctatct	cagcgatctg	tctatctcgt	tcattccatag	ttgcctgact	3480
cccgtcgtg	tagataacta	cgatacggga	gggcttacca	tctggcccca	gtgctgcaat	3540
gataccgcga	gacccaagct	caccggctcc	agatttatca	gcaataaacc	agccagccgg	3600
aagggccgag	cgcagaagtg	gtcctgcaac	tttatccgcc	tccatccagt	ctattaattg	3660
ttgcccggaa	gctagagtaa	gtagttcgcc	agttaatagt	ttgcgcaacg	ttggtggcat	3720
tgctacagcg	atcgtgggtg	cacgctcgtc	gtttggtatg	gcttcattca	gctccggttc	3780
ccaacgatca	aggcgagtta	catgatcccc	catgttgtgc	aaaaaagcgg	ttagctcctt	3840
cggctcctcg	atcgttgtca	gaagtaagtt	ggccgcagtg	ttatcactca	tggttatggc	3900
agcactgcat	aattctctta	ctgtcatgcc	atccgtaaga	tgcttttctg	tgactggtga	3960
gtactcaacc	aagtcattct	gagaataccg	cgccggcgca	ccgagttgct	cttgcccggc	4020
gtcaatacgg	gataatagtg	tatgacatag	cagaacttta	aaagtgtca	tcattggaaa	4080
acgttcttcg	ggcgaaaaac	tctcaaggat	cttaccgctg	ttgagatoca	gttcgatgta	4140
accactcgt	gcaccaact	gatcttcagc	atcttttact	ttcaccagcg	tttctgggtg	4200
agcaaaaaaca	ggaaggcaaa	atgccgcaa	aaagggcaata	aggcgacac	ggaaatggtg	4260
aatactcata	ctcttcttt	ttcaatatta	ttgaagcatt	tatcagggtt	attgtctcat	4320
gagcggatac	atatttgaat	gtatttagaa	aaataaacia	ataggggttc	cgcgcacatt	4380
tcccgaaaa	gtgccacctg	acgtctaaga	aaccattatt	atcatgacat	taacctataa	4440
aaataggcgt	atcacgaggc	cctttcgtct	cgcgcgtttc	ggtgatgacg	gtgaaaacct	4500
ctgacacatg	cagctcccgg	agacggtcac	agcttgtctg	taagcggatg	ccgggagcag	4560
acaagcccgt	cagggcgctg	cagcgggtgt	tggcgggtgt	cgggctggc	ttaactatgc	4620
ggcatcagag	cagattgtac	tga				4643

<210> 5
 <211> 4454
 <212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Plasmid pGX104

<400> 5

```

gagtgcacca tatgcggtgt gaaataccgc acagatgcgt aaggagaaaa taccgcatca 60
ggcgaaattg taaacgttaa tattttgtta aaattcgcgt taaatatttg ttaaatacagc 120
tcatttttta accaataggc cgaaatcggc aaaatccctt ataaatcaaa agaatagacc 180
gagatagggt tgagtgttgt tccagtttgg aacaagagtc cactattaaa gaacgtggac 240
tccaacgtca aagggcgaaa aaccgtctat cagggcgatg gccactacg tgaacatca 300
cccaaatcaa gttttttgcg gtcgaggtgc cgtaaagctc taaatcgaa ccctaaaggg 360
agccccgat ttagagcttg acggggaag ccggcgaacg tggcgagaaa ggaaggaag 420
aaagcgaaag gagcggtcgc tagggcgctg gcaagtgtag cggtcacgct gcgcgtaacc 480
accacaccgc ccgcgcttaa tgcgcgcta cagggcgctg ccattcgcca ttcaggctgc 540
gcaactgttg ggaaggcga tcggtgcggg cctcttcgct attacgccag ctggcgaaag 600
ggggatgtgc tgcaaggcga ttaagttggg taacgccagg gttttccag tcacgacgtt 660
gtaaacgac ggccagtga ttgtaatac actcaactata gggcgaatc aaaaaacccc 720
tcaagaccgc tttagaggcc ccaaggggtt atgctagtga attctgcagg gtaccgggg 780
atcctctaga gatccctcga cctcgagatc cattgtgctg gaccgtggta ctcttatgga 840
gctcggaaatc tcgccaatcg tcactttctg acttatcatg caacttctcg ccggagccaa 900
gatcatcgaa tcgagagaca caccaaagga ccgtgctctt ttcaacggag cccagaaatg 960
taagccgaaa agtgtgtgtt ttcaatctct aatttttgaa cttttcagtg ttcggtatgg 1020
tcatcactgt tggacaagct attgtctacg tcatgtccgg actctacgga gagccatcgg 1080
aaatcgagc tggaatctgt ctcttatcgc tgcctcaact cgttattgcc ggtctcatcg 1140
tctctcttct cgacgagctt ctccaaaagg gatatggtct cggatccgga atttctctct 1200
tcattgccac caacatctgt gaaaccattg tctggaaggc attctccccg gcaacaatga 1260
acaccggagc tggaaaccgag ttcgaaggag ccgtcattgc tcttttccat cttcttgcca 1320
cccgtccga caaggtccgt gcccttcgtg aggctttcta ccgtcaaac ctccaaact 1380
tgatgaactt gatgctact ttccctcgtt ttgcggtggt tatctacttc caaggattcc 1440
gtgtcgacct cccaatcaag tctgcccgtc accgtggaca atacagcagc taccaatca 1500
agctcttcta cacctccaac attccaatca tcttcaatc tgctctcgtc tccaacctct 1560
acgttatctc tcaggtttgt tgcattctcag tagtaccgtt agatgtttat ctttctctag 1620
agggtcaagt tggccgagaa attttttgag ttcattctca agtctgatgg aaaatgttta 1680
ttttcagat gctcgcggga aagttcggag gaaacttctt catcaacctt ctcggtacct 1740
ggtccgataa caccggatac agaagctacc caactggagg actctgctac tatctttcac 1800
caccagagtc ctttggacac atcttcgaag acccaatcca ctgcaccagc acaatggatc 1860
tcgagggatc ttccatacct accagttctg cgctgcagg tcgcgccgcg gactctctag 1920
acgcgtaagc ttactagcat aacccttgg ggcctctaaa cgggtcttga ggggtttttt 1980
gagcttctcg ccctatagtg agtcgtatta cagcttgagt attctatagt gtcacctaaa 2040
tagcttgcg taatcatggt catagctggt tcctgtgtga aattgttatc cgctcacaat 2100
tccacacaac atacgagccg gaagcataaa gtgtaaagcc tgggggtgctt aatgagtgag 2160
ctaactcaca ttaattgctg tgcgctcact gccgccttc cagtcgggaa acctgtcgtg 2220
ccagctgcat taatgaatcg gccaacgcgc ggggagagge ggtttgcgta ttgggcgctc 2280
ttcgccttcc tcgctcactg actcgcctgcg ctcggtcgtt cggctgcggc gagcggatc 2340
agctcactca aaggcggtaa tacggttatc cacagaatca ggggataacg caggaaagaa 2400
catgtgagca aaaggccagc aaaaggccag gaaccgtaaa aaggccgcgt tgctggcgtt 2460
tttcgatagg ctccgcccc ctgacgagca tcacaaaaat cgacgctcaa gtcagaggtg 2520
gcgaaaccgc acaggactat aaagatacca ggcgtttccc cctggaagct cctcgtgcg 2580
ctctcctgtt ccgaccctgc cgttaccgg atacctgtcc gcctttctcc cttcgggaag 2640
cgtggcgtt tctcatagct cacgctgtag gtatctcagt tcggtgtagg tcgttcgctc 2700
caagctgggc tgtgtgcacg aacccccgt tcagccgac cgctgcgctc tatccggtaa 2760
ctatcgtctt gagtccaacc cgtaagaca cgacttatcg ccaactggcag cagcactgg 2820
taacaggatt agcagagcga ggtatgtagg cgggtctaca gagttcttga agtgggtggc 2880
taactacggc tactactagaa ggacagtatt tggatctgct gctctgctga agccagttac 2940
cttcgaaaaa agagttggta gctcttgatc cggcaaaaa accaccgctg gtagcgggtg 3000
ttttttgtt tgcaagcagc agattacgcg cagaaaaaaa ggatctcaag aagatccttt 3060
gatcttttct acggggtctg acgctcagtg gaacgaaaa tcacggttaag ggattttggt 3120

```



```

catgagatta tcaaaaagga tcttcaccta gatcctttta aattaaat gaagttttaa 3180
atcaatctaa agtataatg agtaaacttg gtctgacagt taccaatgct taatcagtga 3240
ggcacctatc tcagcgatct gtctatctcg ttcacccata gtgacctgac tccccgctcg 3300
gtagataact acgatacggg agggcttacc atctggcccc agtgctgcaa tgataccgcg 3360
agaccacgc tcaccgctc cagatttatc agcaataaac cagccagccg gaagggccga 3420
gocgagaagt ggtcctgcaa ctttatccgc ctccatccag tctattaatt gttgccggga 3480
agctagagta agtagttcgc cagttaatag tttgocgaac gttgttgga ttgctacagg 3540
catcgtggtg tcacgctcgt cgtttggtat ggcttcattc agctccggtt cccaacgatc 3600
aagcgaggtt acatgatccc ccattgtgtg caaaaagcg gttagctcct tcggctctcc 3660
gatcgttgtc agaagtaagt tggcgcgagt gttatcactc atggttatgg cagcactgca 3720
taattctctt actgtcatgc catccgtaag atgcttttct gtgactggtg agtactcaac 3780
caagtcattc tgagaatacc gogcccgcg accgagttgc tcttgccggg cgtcaatacg 3840
ggataatagt gtatgacata gcagaacttt aaaagtgtc atcattggaa aacgttcttc 3900
ggggcgaaaa ctctcaagga tcttaccgct gttgagatcc agttcgatgt aaccactcg 3960
tgacccaac tgatcttcag catcttttac tttcaccagc gtttctgggt gagcaaaaac 4020
aggaaggaa aatgccgcaa aaaagggat aagggcgaca cggaaatgtt gaatactcat 4080
actcttctt tttcaatatt attgaagcat ttatcagggt tattgtctca tgagcggata 4140
catatttgaa tgtatttaga aaaataaaca aataggggtt ccgcgacat ttccccgaaa 4200
agtgccacct gacgtctaag aaaccattat tatcatgaca ttaacctata aaaataggcg 4260
tatcacgagg cccttctgct tcgcgctgtt cggtgatgac ggtgaaaacc tctgacacat 4320
gcagctcccg gagacggtca cagcttgtct gtaagcggat gccgggagca gacaagcccg 4380
tcagggcgcg tcagcgggtg ttggcgggtg tcggggctgg cttactatg cggcatcaga 4440
gcagattgta ctga 4454
    
```

- <210> 6
- <211> 4701
- <212> DNA
- <213> Artificial Sequence

- <220>
- <223> Description of Artificial Sequence: Plasmid pGZ18

```

<400> 6
accagcttt cttgtacaaa gtggtgatct ttccagcaca atggatctcg agggatcttc 60
catacctacc agttctgctg ctgcaggctcg cggcgcgac tctctagacg cgtaagctta 120
ctagcataac cccttggggc ctctaaacgg gtcttgaggg gtttttgag cttctcgccc 180
tatagtgagt cgtattacag cttagagtatt ctatagtgtc acctaaatag cttggcgtaa 240
tcatggtcat agctgtttcc tgtgtgaaat tgttatccgc tcacaattcc acacaacata 300
cgagccggaa gcataaagt taaagcctgg ggtgcctaag gagtgagcta actcacatta 360
attgcgttgc gctcactgcc cgctttccag tcgggaaacc tgcctgcca gctgcattaa 420
tgaatcggcc aacgcgcggg gagaggcggg ttgctgattg ggcgctcttc cgcttctcgc 480
ctcactgact cgctgcgctc ggtcgttcgg ctgcggcgag cggtatcagc tactcaaaag 540
gocgtaatac ggttatccac agaatcaggg gataacgcag gaaagaacat gtgagcaaaa 600
ggccagcaaa aggccaggaa ccgtaaaaag gccgcggttc tggcgttttt cgataggctc 660
cgccccctg acgagcatca caaaaatcga cgctcaagtc agaggtggcg aaaccgcaga 720
ggactataaa gataccaggc gtttccccct ggaagctccc tcgtgcgctc tctgttccg 780
accctgccgc ttaccggata cctgtcgcct ttctccctt cgggaagcgt ggcgctttct 840
catagctcac gctgtaggta tctcagttcg gtgtaggtcg ttcgctccaa gctgggctgt 900
gtgcacgaac cccccgttca gcccgaccgc tgcgccttat ccgtaacta tcgtcttgag 960
tccaaccggy taagacacga cttatcgcca ctggcagcag ccaactggtaa caggattagc 1020
agagcgaggt atgtaggcgg tgctacagag ttcttgaagt ggtggcctaa ctacggctac 1080
actagaagga cagtatttgg tatctgcgct ctgctgaagc cagtacctt cggaaaaaga 1140
gttggtagct cttgatccgg caaacaacc accgctggta gcggtgggtt ttttgttgc 1200
aagcagcaga ttcagcgcag aaaaaagga tctcaagaag atcctttgat cttttctacg 1260
gggtctgacg ctcagtggaa cgaaaactca cgttaaggga ttttggctat gagattatca 1320
aaaagatct tcacctagat ctttttaaat taaaaatgaa gttttaaact aaactaaagc 1380
atatatgagt aaacttggtc tgacagttac caatgcttaa tcagtaggc acctatctca 1440
gcgatctgtc tatttcgttc atccatagtt gcctgactcc ccgtcgtgta gataactacy 1500
    
```

atacgggagg	gcttaccatc	tggccccagt	gctgcaatga	taccgcgaga	cccacgctca	1560
ccggctccag	atztatcagc	aataaaccag	ccagccggaa	gggccgagcg	cagaagtggg	1620
cctgcaactt	tatccgcctc	catccagctc	attaattggt	gccgggaagc	tagagtaagt	1680
agttcgccag	ttaatagttt	gcgcaacggt	gttggcattg	ctacagggat	cgtgggtgtca	1740
cgctcgctgt	ttggtatggc	ttcattcagc	tccggttccc	aacgatcaag	gcgagttaca	1800
tgatccccc	tgttgtgcaa	aaaagcgggt	agctccttcg	gtcctccgat	cgttgtcaga	1860
agtaagtgg	ccgcagtggt	atcactcatg	gttatggcag	cactgcataa	ttctcttact	1920
gtcatgccat	ccgtaagatg	cttttctgtg	actggtgagt	actcaaccaa	gtcattctga	1980
gaataaccg	cccgccgacc	gagttgctct	tgcccggcgt	caatacggga	taatagtgt	2040
tgacatagca	gaactttaaa	agtgtcatc	attggaaaac	gttcttcggg	gcgaaaactc	2100
tcaaggatct	taccgctggt	gagatccagt	tcgatgtaac	ccactcgtgc	acccaactga	2160
tcttcagcat	cttttacttt	caccagcgtt	tctgggtgag	caaaaacagg	aaggcaaaat	2220
gccgcaaaaa	aggyaataag	ggcgacacgg	aaatggtgaa	tactcatact	cttccttttt	2280
caatattatt	gaagcattta	tcagggttat	tgtctcatga	gcggatacat	atltgaaatgt	2340
atitagaaaa	ataaacaat	aggggttccg	cgcacatttc	cccgaaaagt	gccacctgac	2400
gtctaagaaa	ccattattat	catgacatta	acctataaaa	ataggcgtat	cacgaggccc	2460
tttctgtctg	cggtttcgg	tgatgacggt	gaaaacctct	gacacatgca	gctcccgtag	2520
acggtcacag	cttgtctgta	agcggatgcc	gggagcagac	aagcccgtca	gggcgcgtca	2580
gcgggtggtg	gcgggtgctg	gggctggcct	aactatgcgg	catcagagca	gattgtactg	2640
agagtgcacc	atatgcggtg	tgaaataccg	cacagatgcg	taaggagaaa	ataccgcatc	2700
aggcgaaatt	gtaaacggtt	atattttggt	aaaattcgcg	ttaaatattt	gttaaatcag	2760
ctcatttttt	aaccaatagg	ccgaaatcgg	caaaatccct	tataaatcaa	aagaatagac	2820
cgagatagg	ttgagtgttg	ttccagtttg	gaacaagagt	ccactattaa	agaacgtgga	2880
ctccaacgtc	aaagggcgaa	aaaccgtcta	tcagggcgat	ggcccactac	gtgaaccata	2940
acccaaatca	agttttttgc	ggtcagagtg	ccgtaaagct	ctaaatcgga	accctaaagg	3000
gagccccga	tttagagctt	gacggggaaa	gccggcgaac	gtggcgagaa	aggaagggaa	3060
gaaagcgaaa	ggagcggcg	ctagggcgct	ggcaagtgt	gcggtcacgc	tgccgcgtaac	3120
caccacaccc	gccgcgctta	atgcgcccgt	acagggcgcg	tccattcgcc	attcaggctg	3180
cgcaactggt	gggaagggcg	atcgggtcgg	gcctcttcgc	tattacgcca	gctggcgaaa	3240
gggggatgtg	ctgcaaggcg	attaagttgg	gtaacgccag	ggttttccca	gtcacgacgt	3300
tgtaaacgca	cgccagtgta	attgtaatac	gactcactat	agggcgaatt	caaaaaaccc	3360
ctcaagacco	gttttagagg	occaaagggt	tatgctagtg	aattctgcag	ggtaccgggt	3420
gatcctctag	agatccctcg	acctcgagat	ccattgtgct	ggaaaacgct	tgcaaggctg	3480
gcaagccacg	tttgggtggtg	gcgaccatcc	tccaaaatca	acaagtttgt	acaaaaaagc	3540
aggctatgcc	aagtacatgt	cgattgcgta	cgcttcgta	atgttggtg	tgttagtcgc	3600
taccagcagt	caaattgttc	tcgagagtgc	gtttttacat	tatcccttca	tcctgattac	3660
gacaattttc	agctgttctc	gotcctacat	ctctcttcat	tgtcacaatg	gtcggaatct	3720
tcttctttgc	tgatgtctt	catccaaaag	aattcacgaa	tattatccat	ggtgtcgtat	3780
tcttctctcat	gattccatct	acatattgtg	tctcacttt	atattcgctc	atcaatctca	3840
acgttatcac	gtggggaact	cgtgaagctg	tcgctaaggc	aacgggacaa	aagacgaaaa	3900
aagcgcctat	ggaacaat	atagacagag	tgattgat	tgtaaaaaag	ggattcagat	3960
taatcagttg	tcgggagaag	aaggaacatg	aagagagacg	agagaaaatg	gaaaagaaaa	4020
tgagagaaat	ggagctagcc	ttgagaagta	ttgaggttat	ctttaacttt	agaaatgtga	4080
aattaataat	ttattttcag	agtggtgccg	acgtgaagaa	aattctcgat	gcaacagagg	4140
agaaggagaa	acgtgaagaa	gaaactcaaa	ctgcagat	tccgattgaa	gagaacgtag	4200
agaagactca	aaaagagatt	cagaaggcaa	accgttatgt	gtggatgaca	agtcatagct	4260
tgaaagtgtg	tgaacgagga	aaactgaaaa	gtcgggaaaa	ggttttctgg	aacgagctca	4320
tcaatgcata	tctgaaaccg	atcaagacga	cgccagctga	aatgaaagcc	gtcggcgaag	4380
gattggcttc	tctacgaaat	cagattgctt	tactattct	tctcgttaat	tctcttcttg	4440
ctcttgccat	ctttttgatt	cagaaacaca	aaaatgtgct	cagcatcaag	ttctcgccaa	4500
tcagtaagca	atattaccct	tatggtcaat	tcaaaaaatt	tgtttttttt	ttctagaaaa	4560
cttccgatgg	acgaaaatga	atgagatgac	tggacaatac	gaggaaaccg	atgaaccatt	4620
aaaaatagat	ccacttgtaa	tgggaattgt	tgttttcctt	ctaattattc	tttttgttca	4680
aactctcgga	atgcttctcc	a				4701

<210> 7
 <211> 25
 <212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:
Oligonucleotide primer C04H5.6F

<400> 7

tgctcagaga gtttctcaac gaacc

25

<210> 8

<211> 25

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:Oligonucleotide
primer C04H5.6R

<400> 8

caatgtagt tgctaggacc acctg

25

<210> 9

<211> 20

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:Oligonucleotide
primer K11D9.2bF

<400> 9

cagccgatct ccgtcttgtg

20

<210> 10

<211> 20

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:Oligonucleotide
primer K11D9.2bR

<400> 10

ccgagggcaa gacaacgaag

20

<210> 11

<211> 24

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:Oligonucleotide
primer Y57G11C.15F

<400> 11

accgtggtac tcttatggag ctcg

24

<210> 12

<211> 20
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence:Oligonucleotide
primer Y57G11C.15R

<400> 12
tgcagtggat tgggtcttcg 20

<210> 13
<211> 52
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence:Oligonucleotide
primer T25G3.2F

<400> 13
ggggacaagt ttgtacaaaa aagcaggcta tgccaagtac atgtcgattg cg 52

<210> 14
<211> 52
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence:Oligonucleotide
primer T25G3.2R

<400> 14
ggggaccact ttgtacaaga aagctggggtt ggagaagcat tccgagagtt tg 52

INTERNATIONAL SEARCH REPORT

International Application No
PCT/EP 00/13149

A. CLASSIFICATION OF SUBJECT MATTER IPC 7 C12N15/10 C12N9/22 C07K14/435 C12N15/66 C12N15/70 C12N1/00				
According to International Patent Classification (IPC) or to both national classification and IPC				
B. FIELDS SEARCHED Minimum documentation searched (classification system followed by classification symbols) IPC 7 C12N				
Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched				
Electronic data base consulted during the international search (name of data base and, where practical, search terms used) BIOSIS, EPO-Internal, WPI Data, PAJ				
C. DOCUMENTS CONSIDERED TO BE RELEVANT				
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.		
X	TIMMONS L ET AL: "Specific interference by ingested dsRNA" NATURE, MACMILLAN JOURNALS LTD. LONDON, GB, vol. 395, no. 6705, 29 October 1998 (1998-10-29), page 854 XP002103601 ISSN: 0028-0836 the whole document	1-4, 12-21, 23-25, 28-31, 34-39		
A	FIRE A ET AL: "Potent and specific genetic interference by double-stranded RNA in Caenorhabditis elegans" NATURE, GB, MACMILLAN JOURNALS LTD. LONDON, vol. 391, 19 February 1998 (1998-02-19), pages 806-811, XP002095876 ISSN: 0028-0836 cited in the application the whole document	1-4		
--- -/--				
<input checked="" type="checkbox"/> Further documents are listed in the continuation of box C.				
<input checked="" type="checkbox"/> Patent family members are listed in annex.				
* Special categories of cited documents :				
<table style="width: 100%; border: none;"> <tr> <td style="width: 50%; border: none; vertical-align: top;"> *A* document defining the general state of the art which is not considered to be of particular relevance *E* earlier document but published on or after the international filing date *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified) *O* document referring to an oral disclosure, use, exhibition or other means *P* document published prior to the international filing date but later than the priority date claimed </td> <td style="width: 50%; border: none; vertical-align: top;"> *T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention *X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone *Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art. *&* document member of the same patent family </td> </tr> </table>			*A* document defining the general state of the art which is not considered to be of particular relevance *E* earlier document but published on or after the international filing date *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified) *O* document referring to an oral disclosure, use, exhibition or other means *P* document published prior to the international filing date but later than the priority date claimed	*T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention *X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone *Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art. *&* document member of the same patent family
A document defining the general state of the art which is not considered to be of particular relevance *E* earlier document but published on or after the international filing date *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified) *O* document referring to an oral disclosure, use, exhibition or other means *P* document published prior to the international filing date but later than the priority date claimed	*T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention *X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone *Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art. *&* document member of the same patent family			
Date of the actual completion of the international search <p style="text-align: center;">5 July 2001</p>	Date of mailing of the international search report <p style="text-align: center;">20/07/2001</p>			
Name and mailing address of the ISA European Patent Office, P.B. 5818 Patentlaan 2 NL - 2280 HV Rijswijk Tel. (+31-70) 340-2040, Tx. 31 651 epo nl. Fax: (+31-70) 340-3016	Authorized officer <p style="text-align: center;">Mateo Rosell, A.M.</p>			

INTERNATIONAL SEARCH REPORT

International Application No
PCT/EP 00/13149

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT		
Category	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	SHARP PHILLIP A: "RNAi and double-strand RNA." GENES & DEVELOPMENT, vol. 13, no. 2, 15 January 1999 (1999-01-15), pages 139-141, XP002171268 ISSN: 0890-9369 the whole document	1-4
A	RAY C ET AL: "GUT-SPECIFIC AND DEVELOPMENTAL EXPRESSION OF A CAENORHABDITIS ELEGANS CYSTEINE PROTEASE GENE" MOLECULAR AND BIOCHEMICAL PARASITOLOGY,NL,ELSEVIER SCIENCE PUBLISHERS, AMSTERDAM, vol. 51, 1992, pages 239-249, XP000572340 ISSN: 0166-6851 abstract	2-4,8
A	RAND J B ET AL: "GENETIC PHARMACOLOGY: INTERACTIONS BETWEEN DRUGS AND GENE PRODUCTS IN CAENORHABDITIS ELEGANS" METHODS IN CELL BIOLOGY,LONDON,GB, vol. 84, 1995, pages 187-204, XP000956211 page 190, paragraph 1 -page 194, paragraph 4	1-4,8
A	AVERY LEON ET AL: "The Caenorhabditis elegans unc-31 gene affects multiple nervous system-controlled functions." GENETICS, vol. 134, no. 2, 1993, pages 455-464, XP001011453 ISSN: 0016-6731 the whole document	2-4,8
A	TAGESSON C ET AL: "INFLUENCE OF SURFACE-ACTIVE FOOD ADDITIVES ON THE INTEGRITY AND PERMEABILITY OF RAT INTESTINAL MUCOSA" FOOD AND CHEMICAL TOXICOLOGY, vol. 22, no. 11, 1984, pages 861-864, XP001009621 ISSN: 0278-6915 the whole document	8

-/--

INTERNATIONAL SEARCH REPORT

International Application No

PCT/EP 00/13149

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT		
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
P,A	<p>LYON CHRISTOPHER J ET AL: "The C. elegans apoptotic nuclease NUC-1 is related in sequence and activity to mammalian DNase II."</p> <p>GENE (AMSTERDAM), vol. 252, no. 1-2, 11 July 2000 (2000-07-11), pages 147-154, XP001009494 ISSN: 0378-1119 abstract; figure 3 page 151, right-hand column, paragraph 2 -page 153, right-hand column, paragraph 2</p>	6,7
P,X	<p>WO 00 01846 A (MORTIER KATHERINE ;DEVGEN NV (BE); BOGAERT THIERRY (BE); PLAETINCK) 13 January 2000 (2000-01-13) cited in the application</p> <p>page 8, line 9 -page 10, line 22 page 12, line 14 -page 23, line 2 figures 5,9; examples 1-4</p>	1-4, 12-21, 23-25, 28-31, 34-39
P,X	<p>WO 00 63425 A (FEICHTINGER RICHARD ;BEGHYN MYRIAM (BE); DEVGEN NV (BE); BOGAERT T) 26 October 2000 (2000-10-26) abstract page 2, line 14-30 page 6, line 18-33 page 7, line 12 -page 9, line 28 page 12, line 20 -page 13, line 25; example 3</p>	2-5,8,9, 29-33
T	<p>TIMMONS LISA ET AL: "Ingestion of bacterially expressed dsRNAs can produce specific and potent genetic interference in Caenorhabditis elegans."</p> <p>GENE (AMSTERDAM), vol. 263, no. 1-2, 2001, pages 103-112, XP001009512 ISSN: 0378-1119 the whole document</p>	1-4, 12-21, 23-26, 28-31, 34-39

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No PCT/EP 00/13149

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 0001846 A	13-01-2000	AU 4907999 A	24-01-2000
		EP 1093526 A	25-04-2001
		GB 2349885 A	15-11-2000
		NO 20010019 A	05-03-2001
WO 0063425 A	26-10-2000	AU 3984600 A	02-11-2000
		GB 2351152 A	20-12-2000